

DESCRIPTIONHERBICIDAL COMPOSITIONTECHNICAL FIELD

The present invention relates to a herbicidal
5 composition.

BACKGROUND ART

As a result of research and development for a long
time, various kinds of various agrochemicals have been
developed and practically used, and these herbicides have
10 contributed to improvement in productivity of
agricultural crops or contributed to elimination or
reduction of labor of removing weeds. However even today,
it is demanded to develop a new agrochemical having a
more satisfactory herbicidal property.

15 It is demanded to provide a herbicidal agent used
for useful crops, which achieves a satisfactory
herbicidal effect at a small dose and has an excellent
selectivity between aimed crops and weeds by being
applied to soil or plant foliage.

20 DISCLOSURE OF THE INVENTION

An isoxazoline compound of the formula (I) which is
one of active ingredients in the herbicidal composition
of the present invention is safe to rice, wheat, barley,
corn, grain sorghum, soybeans, cotton, sugar beet, turf,
25 fruit trees and the like, and has an excellent herbicidal
effect by itself.


The present inventors have discovered that by

combining an isoxazoline derivative of the formula (I) with at least one herbicide of Group A at a predetermined ratio, not a simple total herbicidal effect but a synergistic herbicidal effect can be achieved. Thus, by combining at least two agrochemical agents, a herbicidal spectrum is broadened as compared with a herbicidal application range by each agent, and a herbicidal effect is achieved at an earlier stage and is retained for a longer time, and a satisfactory herbicidal effect can be achieved at a smaller dose than a dose by a single use of each agent. Also, the herbicidal composition prepared by combining at least two agrochemical agents is safe to rice, wheat, barley, corn, grain sorghum, soybeans, cotton, sugar beet, turf, fruit trees and the like, and achieves a satisfactory herbicidal effect by one treatment. The present invention has been accomplished on the basis of this discovery.

The present invention is characterized by having the following features.

1. A herbicidal composition which comprises i) an isoxazoline derivative represented by the following general formula (I) or its salt and ii) at least one compound selected from the following Group A:

i) Formula (I)



wherein R^1 and R^2 are respectively independently a hydrogen atom, a C1 to C10 alkyl group, a C3 to C8 cycloalkyl group or a C3 to C8 cycloalkyl C1 to C3 alkyl group; or R^1 and R^2 may be bonded to each other to form a C3 to C7 spiro ring together with the carbon atoms to which they bond;

R^3 and R^4 are respectively independently a hydrogen atom, a C1 to C10 alkyl group or a C3 to C8 cycloalkyl group; or R^3 and R^4 may be bonded to each other to form a C3 to C7 spiro ring together with the carbon atoms to which they bond; or R^1 , R^2 , R^3 and R^4 may form a 5- to 8-membered ring together with the carbon atoms to which they bond;

R^5 and R^6 are respectively independently a hydrogen atom or a C1 to C10 alkyl group;

Y is a 5- to 6-membered aromatic heterocyclic group or condensed aromatic heterocyclic group having one or more hetero atoms selected from a nitrogen atom, an oxygen atom and a sulfur atom; the heterocyclic group may be substituted with 0 to 6 same or different groups selected from the following substituent group α ; when the heterocyclic group is substituted at the two adjacent

positions with two alkyl groups, two alkoxy groups, an alkyl group and an alkoxy group, an alkyl group and an alkylthio group, an alkyl group and an alkylsulfonyl group, an alkyl group and a monoalkylamino group, or an
 5 alkyl group and a dialkylamino group, all selected from the substituent group α , the two groups may form, together with the atoms to which they bond, a 5- to 8-membered ring which may be substituted with 1 to 4 halogen atoms; the hetero atom of the heterocyclic group,
 10 when it is a nitrogen atom, may be oxidized to become N-oxide;

n is an integer of 0 to 2;

(Substituent group α)

Hydroxyl group; thiol group; halogen atoms; C1 to
 15 C10 alkyl groups; C1 to C10 alkyl groups each mono-substituted with a group selected from the following substituent group β , C1 to C4 haloalkyl groups; C3 to C8 cycloalkyl groups; C1 to C10 alkoxy groups; C1 to C10 alkoxy groups each mono-substituted with a group selected
 20 from the following substituent group γ ; C1 to C4 haloalkoxy groups; C3 to C8 cycloalkyloxy groups; C3 to C8 cycloalkyl C1 to C3 alkyloxy groups; C1 to C10 alkylthio groups; C1 to C10 alkylthio groups each mono-substituted with a group selected from the substituent
 25 group γ ; C1 to C4 haloalkylthio groups; C2 to C6 alkenyl groups; C2 to C6 alkenyloxy groups; C2 to C6 alkynyl groups; C2 to C6 alkynyloxy groups; C1 to C10

alkylsulfinyl groups; C1 to C10 alkylsulfinyl groups each mono-substituted with a group selected from the substituent group γ ; C1 to C10 alkylsulfonyl groups; C1 to C10 alkylsulfonyl groups each mono-substituted with a group selected from the substituent group γ ; C1 to C4 haloalkylsulfinyl groups; C1 to C10 alkylsulfonyloxy groups each mono-substituted with a group selected from the substituent group γ ; C1 to C4 haloalkylsulfonyl groups; C1 to C10 alkylsulfonyloxy groups; C1 to C4 haloalkylsulfonyloxy groups; optionally substituted phenyl group; optionally substituted phenoxy group; optionally substituted phenylthio group; optionally substituted aromatic heterocyclic groups; optionally substituted aromatic heterocyclic oxy groups; optionally substituted aromatic heterocyclic thio groups; optionally substituted phenylsulfinyl groups; optionally substituted phenylsulfonyl groups; optionally substituted aromatic heterocyclic sulfonyl groups; optionally substituted phenylsulfonyloxy groups; acyl groups; C1 to C4 haloalkylcarbonyl groups; optionally substituted benzylcarbonyl group; optionally substituted benzoyl group; carboxyl group; C1 to C10 alkoxy carbonyl groups; optionally substituted benzyloxycarbonyl group; optionally substituted phenoxycarbonyl group; cyano group; carbamoyl group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups and optionally substituted phenyl

group); C1 to C6 acyloxy groups; C1 to C4 haloalkylcarbonyloxy groups; optionally substituted benzylcarbonyloxy group; optionally substituted benzoyloxy group; nitro group; and amino group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups, optionally substituted phenyl group, C1 to C6 acyl groups, C1 to C4 haloalkylcarbonyl groups, optionally substituted benzylcarbonyl group, optionally substituted benzoyl group, C1 to C10 alkylsulfonyl group, C1 to C4 haloalkylsulfonyl groups, optionally substituted benzylsulfonyl group, and optionally substituted phenylsulfonyl group);

(Substituent group β)

Hydroxyl group; C3 to C8 cycloalkyl groups (which may be substituted with halogen atom or alkyl group); C1 to C10 alkoxy groups; C1 to C10 alkylthio groups; C1 to C10 alkylsulfonyl groups; C1 to C10 alkoxycarbonyl groups; C2 to C6 haloalkenyl groups; amino group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups, C1 to C6 acyl groups; C1 to C4 haloalkylcarbonyl groups, C1 to C10 alkylsulfonyl groups and C1 to C4 haloalkylsulfonyl groups); carbamoyl group (its nitrogen atom may be substituted with same or different C1 to C10 alkyl groups); C1 to C6 acyl groups; C1 to C4 haloalkylcarbonyl groups; C1 to C10 alkoxyimino groups; cyano group;

optionally substituted phenyl group; and optionally substituted phenoxy group;

(Substituent group γ)

C1 to C10 alkoxy carbonyl groups; optionally substituted phenyl group; optionally substituted aromatic heterocyclic groups; cyano group; and carbamoyl group (its nitrogen atom may be substituted with same or different C1 to C10 alkyl groups); and

ii) Group A

atrazine, simazine, cyanazine, isoxaflutole, mesotrione, flumetsulam, imazethapyr, imazapyr, dicamba, clopyralid, prosulfuron, halosulfuron-methyl, rimsulfuron, bentazone, carfentrazone-ethyl, metribuzin, thifensulfuron-methyl, nicosulfuron, primisulfuron, cloransulam-methyl, glufosinate, glyphosate, glyphosate-trimesium, pendimethalin, linuron, prometryn, diflufenican, flumioxazin, and metolachlor.

2. The herbicidal composition according to Feature 1, wherein the isoxazoline derivative of the formula (I) or its salt has a substituent selected from the substituent group α on the heterocycle which may be substituted with 0 to 6 same or different groups, including hydroxyl group; halogen atoms; C1 to C10 alkyl groups; C1 to C10 alkyl groups each mono-substituted with a group selected from the substituent group β , C1 to C4 haloalkyl groups; C3 to C8 cycloalkyl groups; C1 to C10 alkoxy groups; C1 to C10 alkoxy groups each mono-substituted with a group

selected from the substituent group γ ; C1 to C4
 haloalkoxy groups; C3 to C8 cycloalkyloxy groups; C3 to
 C8 cycloalkyl C1 to C3 alkyloxy groups; C1 to C10
 alkylthio groups; C1 to C10 alkylthio groups each mono-
 5 substituted with a group selected from the substituent
 group γ ; C1 to C4 haloalkylthio groups; C2 to C6 alkenyl
 groups; C2 to C6 alkenyloxy groups; C2 to C6 alkynyl
 groups; C2 to C6 alkynyloxy groups; C1 to C10
 alkylsulfonyl groups; C1 to C4 haloalkylsulfonyl groups;
 10 optionally substituted phenyl group; optionally
 substituted phenoxy group; optionally substituted
 phenylthio group; optionally substituted aromatic
 heterocyclic groups; optionally substituted aromatic
 heterocyclic oxy groups; optionally substituted aromatic
 15 heterocyclic thio groups; optionally substituted
 phenylsulfonyl groups; optionally substituted aromatic
 heterocyclic sulfonyl groups; C1 to C6 acyl groups; C1 to
 C4 haloalkylcarbonyl groups; optionally substituted
 benzylcarbonyl group; optionally substituted benzoyl
 20 group; carboxyl group; C1 to C10 alkoxycarbonyl groups;
 cyano group; carbamoyl group (its nitrogen atom may be
 substituted with same or different groups selected from
 C1 to C10 alkyl groups and optionally substituted phenyl
 group); nitro group; and amino group (its nitrogen atom
 25 may be substituted with same or different groups selected
 from C1 to C10 alkyl groups, optionally substituted
 phenyl group, C1 to C6 acyl groups, C1 to C4

haloalkylcarbonyl groups, optionally substituted
benzylcarbonyl group, optionally substituted benzoyl
group, C1 to C10 alkylsulfonyl groups, C1 to C4
haloalkylsulfonyl groups, optionally substituted
5 benzylsulfonyl group, and optionally substituted
phenylsulfonyl group); when the heterocyclic group is
substituted at the two adjacent positions with two alkyl
groups, two alkoxy groups, an alkyl group and an alkoxy
group, an alkyl group and an alkylthio group, an alkyl
10 group and an alkylsulfonyl group, an alkyl group and a
monoalkylamino group, or an alkyl group and a
dialkylamino group, the two groups may form, together
with the atoms to which they bond, a 5- to 8-membered
ring which may be substituted with 1 to 4 halogen atoms.

15 3. The herbicidal composition according to Feature 2,
wherein the isoxazoline derivative of the formula (I) or
its salt has a substituent selected from the substituent
group α on the heterocycle which may be substituted with
0 to 6 same or different groups, including halogen atoms;
20 C1 to C10 alkyl groups; C1 to C4 haloalkyl groups; C1 to
C10 alkoxy C1 to C3 alkyl groups; C3 to C8 cycloalkyl
groups (which may be substituted with halogen atom or
alkyl group); C1 to C10 alkoxy groups; C1 to C4
haloalkoxy groups; C3 to C8 cycloalkyl C1 to C3 alkyloxy
25 groups; optionally substituted phenoxy group; C1 to C10
alkylthio groups; C1 to C10 alkylsulfonyl groups; acyl
groups; C1 to C4 haloalkylcarbonyl groups; C1 to C10

alkoxycarbonyl groups; cyano group and carbamoyl group (its nitrogen atom may be substituted with same or different C1 to C10 alkyl groups).

4. The herbicidal composition according to any of
5 Feature 1, 2 or 3, wherein R^1 and R^2 may be the same or different and are each a methyl group or an ethyl group; and R^3 , R^4 , R^5 and R^6 are each a hydrogen atom.

5. The herbicidal composition according to any of
Feature 1, 2, 3 or 4, wherein Y is a 5- or 6-membered
10 aromatic heterocyclic group having a hetero atom selected from a nitrogen atom, an oxygen atom and a sulfur atom.

6. The herbicidal composition according to Feature 5, wherein Y is a thienyl group, a pyrazolyl group, an isoxazolyl group, an isothiazolyl group, a pyridyl group
15 or a pyrimidinyl group.

7. The herbicidal composition according to Feature 6, wherein Y is a thiophen-3-yl group, a pyrazol-4-yl group, a pyrazol-5-yl group, an isoxazol-4-yl group, an isothiazol-4-yl group, a pyridyn-3-yl group or a
20 pyrimidin-5-yl group.

8. The herbicidal composition according to Feature 7, wherein Y is a thiophen-3-yl group and the thiophene ring is substituted with the substituent group α at the 2- and 4-positions.

25 9. The herbicidal composition according to Feature 7, wherein Y is a pyrazol-4-yl group and the pyrazole ring is substituted at the 3- and 5-positions with the

substituent group α and at the 1-position with a hydrogen atom, a C1 to C10 alkyl group, a C1 to C10 alkyl group mono-substituted with a group selected from the substituent group β , a C1 to C4 haloalkyl group, a C3 to C8 cycloalkyl group, a C2 to C6 alkenyl group, a C2 to C6 alkynyl group, a C1 to C10 alkylsulfinyl group, a C1 to C10 alkylsulfonyl group, a C1 to C10 alkylsulfonyl group mono-substituted with a group selected from the substituent group γ , a C1 to C4 haloalkylsulfonyl group, an optionally substituted phenyl group, an optionally substituted aromatic heterocyclic group, an optionally substituted phenylsulfonyl group, an optionally substituted aromatic heterocyclic sulfonyl group, an acyl group, a C1 to C4 haloalkylcarbonyl group, an optionally substituted benzylcarbonyl group, an optionally substituted benzoyl group, a C1 to C10 alkoxy carbonyl group, an optionally substituted benzyloxy carbonyl group, an optionally substituted phenoxy carbonyl group, a carbamoyl group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups and optionally substituted phenyl group), or an amino group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups, an optionally substituted phenyl group, acyl groups, C1 to C4 haloalkylcarbonyl groups, an optionally substituted benzylcarbonyl group, an optionally substituted benzoyl group, C1 to C10 alkylsulfonyl groups,

C1 to C4 haloalkylsulfonyl groups, an optionally substituted benzylsulfonyl group and an optionally substituted phenylsulfonyl group).

10. The herbicidal composition according to Feature 7,
5 wherein Y is a pyrazol-5-yl group and the pyrazole ring is substituted at the 4-position with the substituent group α and at the 1-position with a hydrogen atom, a C1 to C10 alkyl group, a C1 to C10 alkyl group mono-substituted with a group selected from the substituent
10 group β , a C1 to C4 haloalkyl group, a C3 to C8 cycloalkyl group, a C2 to C6 alkenyl group, a C2 to C6 alkynyl group, a C1 to C10 alkylsulfinyl group, a C1 to C10 alkylsulfonyl group, a C1 to C10 alkylsulfonyl group mono-substituted with a group selected from the
15 substituent group γ , a C1 to C4 haloalkylsulfonyl group, an optionally substituted phenyl group, an optionally substituted aromatic heterocyclic group, an optionally substituted phenylsulfonyl group, an optionally substituted aromatic heterocyclic sulfonyl group, an acyl
20 group, a C1 to C4 haloalkylcarbonyl group, an optionally substituted benzylcarbonyl group, an optionally substituted benzoyl group, a C1 to C10 alkoxy carbonyl group, an optionally substituted benzyloxycarbonyl group, an optionally substituted phenoxycarbonyl group, a
25 carbamoyl group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups and an optionally substituted phenyl group),

or an amino group (its nitrogen atom may be substituted with same or different groups selected from C1 to C10 alkyl groups, an optionally substituted phenyl group, acyl groups, C1 to C4 haloalkylcarbonyl groups, an optionally substituted benzylcarbonyl group, an optionally substituted benzoyl group, C1 to C10 alkylsulfonyl groups, C1 to C4 haloalkylsulfonyl groups, an optionally substituted benzylsulfonyl group and an optionally substituted phenylsulfonyl group).

11. The herbicidal composition according to Feature 7, wherein Y is an isoxazol-4-yl group and the isoxazole ring is substituted with the substituent group α at the 3- and 5-positions.

12. The herbicidal composition according to Feature 7, wherein Y is an isothiazol-4-yl group and the isothiazole ring is substituted with the substituent group α at the 3- and 5-positions.

13. The herbicidal composition according to Feature 7, wherein Y is a pyridin-3-yl group and the pyridine ring is substituted with the substituent group α at the 2- and 4-positions.

14. The herbicidal composition according to Feature 7, wherein Y is a pyrimidin-5-yl group and the pyrimidine ring is substituted with the substituent group α at the 4- and 6-positions.

15. The herbicidal composition according to any of Features 1 to 14, wherein n is an integer of 2.

16. The herbicidal composition according to any of Features 1 to 14, wherein the compound of Group A is at least one compound selected from the group consisting of atrazine, cyanazine, simazine and prometryn.

5 17. The herbicidal composition according to any of Features 1 to 14, wherein the compound of Group A is at least one compound selected from the group consisting of glyphosate, glufosinate, linuron and flumetsulam.

18. A herbicidal composition which comprises i) the
10 isoxazoline derivative or its salt is a compound as defined in Feature 9 and ii) the compound of Group A is at least one compound selected from the group consisting of atrazine, cyanazine, simazine, prometryn, glyphosate, glufosinate, linuron, flumetsulam, metribuzin,
15 isoxaflutole, mesotrione, diflufenican, pendimethalin and flumioxazin.

19. A herbicidal composition which comprises i) the isoxazoline derivative or its salt is a compound as defined in Feature 9 and ii) the compound of Group A is
20 at least one compound selected from the group consisting of atrazine, cyanazine, simazine and prometryn.

20. A herbicidal composition which comprises i) the isoxazoline derivative or its salt is a compound as defined in Feature 9 and ii) the compound of Group A is
25 at least one compound selected from the group consisting of glyphosate, glufosinate, linuron and flumetsulam.

21. The herbicidal composition according to any of

Features 1 to 20, wherein ii) at least one compound of Group A is contained in an amount of from 0.001 to 100 parts by weight to 1 part by weight of i) an isoxazoline derivative represented by the Formula (I) or its salt.

- 5 22. The herbicidal composition according to any of Features 1 to 21, which is used as an agrochemical product containing i) an isoxazoline derivative of the Formula (I) or its salt and ii) at least one compound of Group A in a total amount of from 0.5 to 90 wt%.

10 The definitions of the terms used in the present specification are given below.

The expression of "C1 to C10", etc. indicates that the substituent appearing after the expression has 1 to 10 carbon atoms in the case of "C1 to C10".

- 15 Halogen atom refers to a fluorine atom, a chlorine atom, a bromine atom or an iodine atom.

C1 to C10 alkyl group refers to a straight or branched chain alkyl group of 1 to 10 carbon atoms unless other wise specified; and there can be mentioned, for
20 example, methyl group, ethyl group, n-propyl group, isopropyl group, n-butyl group, isobutyl group, sec-butyl group, tert-butyl group, n-pentyl group, isopentyl group, neopentyl group, n-hexyl group, isohexyl group, 3,3-dimethylbutyl group, heptyl group and octyl group.

- 25 C3 to C8 cycloalkyl group refers to a cycloalkyl group of 3 to 8 carbon atoms; and there can be mentioned, for example, cyclopropyl group, cyclobutyl group,

cyclopentyl group and cyclohexyl group.

C3 to C8 cycloalkyl C1 to C3 alkyl group (which may be substituted with halogen atom or alkyl group) refers, unless otherwise specified, to a C1 to C3 alkyl group substituted with a C3 to C8 cycloalkyl group which may be substituted with 1 to 4 same or different halogen atoms or C1 to C3 alkyl group; and there can be mentioned, for example, cyclopropylmethyl group, 1-cyclopropylethyl group, 2-cyclopropylethyl group, 1-cyclopropylpropyl group, 2-cyclopropylpropyl group, 3-cyclopropylpropyl group, cyclobutylmethyl group, cyclopentylmethyl group, cyclohexylmethyl group, 2-chlorocyclopropylmethyl group, 2,2-dichlorocyclopropylmethyl group, 2-fluorocyclopropylmethyl group, 2,2-difluorocyclopropylmethyl group, 2-methylcyclopropylmethyl group, 2,2-dimethylcyclopropylmethyl group and 2-methylcyclopropylethyl group.

C3 to C8 cycloalkyl C1 to C3 alkyl group refers to a alkyl group of 1 to 3 carbon atoms, substituted with a cycloalkyl group of 3 to 8 carbon atoms; and there can be mentioned, for example, cyclopropylmethyl group, 1-cyclopropylethyl group, 2-cyclopropylethyl group, 1-cyclopropylpropyl group, 2-cyclopropylpropyl group, 3-cyclopropylpropyl group, cyclobutylmethyl group, cyclopentylmethyl group and cyclohexylmethyl group.

C1 to C4 haloalkyl group refers, unless otherwise

specified, to a straight or branched chain alkyl group of 1 to 4 carbon atoms, substituted with 1 to 9 same or different halogen atoms; and there can be mentioned, for example, fluoromethyl group, chloromethyl group, bromomethyl group, difluoromethyl group, trifluoromethyl group, 2,2-difluoroethyl group, 2,2,2-trifluoroethyl group and pentafluoroethyl group.

C2 to C6 alkenyl group refers to a straight or branched chain alkenyl group of 2 to 6 carbon atoms; and there can be mentioned, for example, ethenyl group, 1-propenyl group, 2-propenyl group, isopropenyl group, 1-butenyl group, 2-butenyl group, 3-butenyl group and 2-pentenyl group.

C2 to C6 alkynyl group refers to a straight or branched chain alkynyl group of 2 to 6 carbon atoms; and there can be mentioned, for example, ethynyl group, 2-propynyl group, 1-methyl-2-propynyl group, 2-butyne group, 3-butyne group and 2-methyl-3-butyne group.

C2 to C6 haloalkenyl group refers, unless otherwise specified, to a straight or branched alkenyl group of 2 to 6 carbon atoms, substituted with 1 to 4 same or different halogen atoms; and there can be mentioned, for example, 3-chloro-2-propenyl group and 2-chloro-2-propenyl group.

C1 to C10 alkoxy group refers to an (alkyl)-O-group wherein the alkyl moiety has the above definition; and there can be mentioned, for example, methoxy group,

ethoxy group, n-propoxy group, isopropoxy group, tert-butoxy group, n-butoxy group, sec-butoxy group and isobutoxy group.

C1 to C4 haloalkoxy group refers to a (haloalkyl)-
 5 O- group wherein the haloalkyl moiety has the above definition; and there can be mentioned, for example, difluoromethoxy group, trifluoromethoxy group, 2,2-difluoroethoxy group and 2,2,2-trifluoroethoxy group.

C3 to C8 cycloalkyloxy group refers to a
 10 (cycloalkyl)-O- group wherein the cycloalkyl moiety has the above definition; and there can be mentioned, for example, cyclopropyloxy group, cyclobutyloxy group, cyclopentyloxy group and cyclohexyloxy group.

C3 to C8 cycloalkyl C1 to C3 alkyloxy group refers
 15 to a (cycloalkylalkyl)-O- group wherein the cycloalkylalkyl moiety has the above definition; and there can be mentioned, for example, cyclopropylmethoxy group, 1-cyclopropylethoxy group, 2-cyclopropylethoxy group, 1-cyclopropylpropoxy group, 2-cyclopropylpropoxy group, 3-cyclopropylpropoxy group, cyclobutylmethoxy group, cyclopentylmethoxy group and cyclohexylmethoxy group.

C2 to C6 alkenyloxy group and C2 to C6 alkynyloxy group refer, respectively, to an (alkenyl)-O- group and
 25 an (alkynyl)-O- group, in each of which the alkenyl or alkynyl moiety has the above definition; and there can be mentioned, for example, 2-propenyloxy group and 2-

propynyloxy group.

C1 to C10 alkoxyimino group refers to an (alkoxy)-N= group wherein the alkoxy moiety has the above definition; and there can be mentioned, for example, methoxyimino
5 group and ethoxyimino group.

C1 to C10 alkylthio group, C1 to C10 alkylsulfinyl group and C1 to C10 alkylsulfonyl group refer, respectively, to an (alkyl)-S- group, an (alkyl)-SO- group and an (alkyl)-SO₂- group, in each of which the
10 alkyl moiety has the above definition; and there can be mentioned, for example, methylthio group, ethylthio group, n-propylthio group, isopropylthio group, methylsulfinyl group, methylsulfonyl group, ethylsulfonyl group, n-propylsulfonyl group and isopropylsulfonyl group.

15 C1 to C10 alkylsulfonyloxy group refers to an (alkylsulfonyl)-O- group wherein the alkylsulfonyl moiety has the above definition, and there can be mentioned, for example, methylsulfonyloxy group and ethylsulfonyloxy group.

20 C1 to C10 alkoxycarbonyl group refers to an (alkoxy)-CO- group wherein the alkoxy moiety has the above definition, and there can be mentioned, for example, methoxycarbonyl group, ethoxycarbonyl group, n-propoxycarbonyl group and isopropoxycarbonyl group.

25 C1 to C6 acryl group refers to a straight or branched chain aliphatic acyl group of 1 to 6 carbon atoms, and there can be mentioned, for example, formyl

group, acetyl group, propionyl group, isopropionyl group, butyryl group and pivaloyl group.

C1 to C10 acyloxy group refers to an (acyl)-O- group wherein the acyl moiety has the above definition; and
5 there can be mentioned, for example, acetoxo group, propionyloxy group, ispropionyloxy group and pivalolyoxy group.

C1 to C4 haloalkylcarbonyl group, C1 to C4 haloalkylthio group and C1 to C4 haloalkylsulfonyl group
10 refers, respectively, to a (haloalkyl)-CO- group, a (haloalkyl)-S- group and a (haloalkyl)-SO₂- group, in each of which the haloalkyl moiety has the above definition; and there can be mentioned, for example, chloroacetyl group, trifluoroacetyl group,
15 pentafluoropropyl group, difluoromethylthio group, trifluoromethylthio group, chloromethylsulfonyl group, difluoromethylsulfonyl group and trifluoromethylsulfonyl group.

C1 to C4 haloalkylcarbonyloxy group and C1 to C4 haloalkylsulfonyloxy group refer, respectively, to a
20 (haloalkylcarbonyl)-O- group and a (haloalkylsulfonyl)-O- group, in each of which the haloalkylcarbonyl moiety or the haloalkylsulfonyl moiety has the above definition; and there can be mentioned, for example, chloroacetyloxy
25 group, trifluoroacetyloxy group, chloromethylsulfonyloxy group and trifluoromethylsulfonyloxy group.

"Optionally substituted" in (optionally

substituted) phenyl group, (optionally substituted)
aromatic heterocyclic group, (optionally substituted)
phenoxy group, (optionally substituted aromatic
heterocyclic oxy group, (optionally substituted)

5 phenylthio group, (optionally substituted) aromatic
heterocyclic thio group, (optionally substituted)
phenylsulfonyl group, (optionally substituted)
phenylsulfonyloxy group, (optionally substituted)
aromatic heterocyclic sulfonyl group, (optionally

10 substituted) benzylcarbonyl group, (optionally
substituted) benzylcarbonyloxy group, (optionally
substituted) benzylsulfonyl group, (optionally
substituted) benzoyl group, (optionally substituted)
benzoyloxy group, (optionally substituted)

15 benzyloxycarbonyl group and (optionally substituted)
phenoxycarbonyl group, refers to being optionally
substituted with, for example, halogen atom, C1 to C10
alkyl group, C1 to C4 haloalkyl group, C1 to C10
alkoxyalkyl group, C1 to C10 alkoxy group, C1 to C10

20 alkylthio group, C1 to C10 alkylsulfonyl group, acyl
group, C1 to C10 alkoxycarbonyl group, cyano group,
carbamoyl group (its nitrogen atom may be substituted
with same or different C1 to C10 alkyl groups), nitro
group or amino group (its nitrogen atom may be

25 substituted with same or different groups selected from
C1 to C10 alkyl groups, C1 to C6 acyl groups, C1 to C4
haloalkylcarbonyl groups, C1 to C10 alkylsulfonyl groups

and C1 to C4 haloalkylsulfonyl groups).

5- to 6-membered aromatic heterocyclic group having a hetero atom selected from a nitrogen atom, an oxygen atom and a sulfur atom includes, for example, furyl group, thienyl group, pyrrolyl group, pyrazolyl group, isoxazolyl group, isothiazolyl group, oxazolyl group, thiazolyl group, imidazolyl group, pyridyl group, pyridazinyl group, pyrimidinyl group, pyrazinyl group, triazinyl group, triazolyl group, oxadiazolyl group and thiadiazolyl group, each having 1 to 3 hetero atoms.

Fused aromatic heterocyclic group refers to a group having 1 to 3 hetero atoms randomly selected from nitrogen atom, oxygen atom and sulfur atom; and there can be mentioned, for example, benzofuryl group, benzothienyl group, indolyl group, benzoxazolyl group, benzothiazolyl group, benzimidazolyl group, benzisoxazolyl group, benzisothiazolyl group, indazolyl group, quinolyl group, isoquinolyl group, phthalazinyl group, quinoxalinyl group, quinazolinyl group, cinnolinyl group and benzotriazolyl group.

Aromatic heterocycle in (optionally substituted) aromatic heterocyclic group, (optionally substituted) aromatic heterocyclic oxy group, (optionally substituted) aromatic heterocyclic thio group and (optionally substituted) aromatic heterocyclic sulfonyl group, refers to a 5- to 6-membered group having 1 to 3 hetero atoms randomly selected from nitrogen atom, oxygen atom and

sulfur atom; and there can be mentioned, for example, furyl group, thienyl group, pyrrolyl group, pyrazolyl group, isoxazolyl group, isothiazolyl group, oxazolyl group, thiazolyl group, imidazolyl group, pyridyl group, pyridazinyl group, pyrimidinyl group, pyrazinyl group, triazinyl group, triazolyl group, oxadiazolyl group and thiadiazolyl group.

Pharmaceutically acceptable salt is a salt of a compound of the general formula [I] having, in the structure, hydroxyl group, carboxyl group, amino group or the like, with a metal or an organic base or with a mineral acid or an organic acid. As the metal, there can be mentioned alkali metals such as sodium, potassium and the like; and alkaline earth metals such as magnesium, calcium and the like. As the organic base, there can be mentioned triethylamine, diisopropylamine, etc. As the mineral acids, there can be mentioned hydrochloric acid, sulfuric acid, etc. As the organic acid, there can be mentioned acetic acid, methanesulfonic acid, p-toluenesulfonic acid, etc.

In the above-mentioned general formula [I], it is preferred that

R^1 and R^2 may be the same or different and are each a methyl group or an ethyl group;

R^3 , R^4 , R^5 and R^6 are each a hydrogen atom;

n is an integer of 2; and

Y is a thiophen-3-yl group [the 2- and 4-positions of the

group are substituted with same or different groups selected from halogen atoms, alkyl groups, haloalkyl groups, alkoxyalkyl groups, cycloalkyl groups, alkoxy groups, haloalkoxy groups, acyl groups, haloalkylcarbonyl groups, alkoxycarbonyl groups, cyano group and carbamoyl group (its nitrogen atom may be substituted with same or different alkyl groups)], or

a pyrazol-4-yl group [the 3- and 5-positions of the group are substituted with same or different groups selected from halogen atoms, alkyl groups, haloalkyl groups, alkoxyalkyl groups, cycloalkyl groups, alkoxy groups, haloalkoxy groups, cycloalkylalkyloxy groups, optionally substituted phenoxy group, alkylthio groups, alkylsulfonyl groups, acyl groups, haloalkylcarbonyl groups, alkoxycarbonyl groups, cyano group and carbamoyl group (its nitrogen atom may be substituted with same or different alkyl groups); the 1-position is substituted with hydrogen atom, alkyl group, alkyl group mono-substituted with a group selected from the substituent group β , haloalkyl group, cycloalkyl group, alkenyl group, alkynyl group, alkylsulfonyl group, alkylsulfonyl group mono-substituted with a group selected from the substituent group γ , haloalkylsulfonyl group, optionally substituted phenyl group, optionally substituted aromatic heterocyclic group, optionally substituted phenylsulfonyl group, optionally substituted aromatic heterocyclicsulfonyl group, acyl group, haloalkylcarbonyl

group, optionally substituted benzylcarbonyl group,
optionally substituted benzoyl group, alkoxycarbonyl
group, optionally substituted benzyloxycarbonyl group,
optionally substituted phenoxycarbonyl group or carbamoyl
5 group (its nitrogen atom may be substituted with same or
different groups selected from alkyl groups and
optionally substituted phenyl group)], or

a pyrazol-5-yl group [the 4-position of the group is
substituted with halogen atom, alkyl group, haloalkyl
10 group, alkoxyalkyl group, haloalkoxy group, acyl group,
haloalkylcarbonyl group, alkoxycarbonyl group, cyano
group or carbamoyl group (its nitrogen atom maybe
substituted with same or different alkyl groups); the 1-
position is substituted with hydrogen atom, alkyl group,
15 alkyl group mono-substituted with a group selected from
the substituent group β , haloalkyl group, cycloalkyl
group, or optionally substituted phenyl group)], or

an isoxazol-4-yl group [the 3- and 5-positions of
the group are substituted with same or different groups
20 selected from halogen atoms, alkyl groups, haloalkyl
groups, alkoxyalkyl groups, cycloalkyl groups, alkoxy
groups, haloalkoxy groups, alkylthio groups,
alkylsulfonyl groups, acyl groups, haloalkylcarbonyl
groups, alkoxycarbonyl groups, cyano group and carbamoyl
25 group (its nitrogen atom may be substituted with same or
different alkyl groups)], or

an isothiazol-4-yl group [the 3- and 5-positions of

the group are substituted with same or different groups selected from halogen atoms, alkyl groups, haloalkyl groups, alkoxyalkyl groups, cycloalkyl groups, alkoxy groups, haloalkoxy groups, optionally substituted phenoxy group, alkylthio groups, alkylsulfonyl groups, acyl groups, haloalkylcarbonyl groups, alkoxycarbonyl groups, cyano group and carbamoyl group (its nitrogen atom may be substituted with same or different alkyl groups)], or

a pyridin-3-yl group [the 2- and 4-positions of the group are substituted with same or different groups selected from halogen atoms, alkyl groups, haloalkyl groups, alkoxyalkyl groups, cycloalkyl groups, alkoxy groups, haloalkoxy groups, alkylthio groups, alkylsulfonyl groups, acyl groups, haloalkylcarbonyl groups, alkoxycarbonyl groups, cyano group and carbamoyl group (its nitrogen atom may be substituted with same or different alkyl groups)], or

a pyrimidin-5-yl group [the 4- and 6-positions of the group are substituted with same or different groups selected from halogen atoms, alkyl groups, haloalkyl groups, alkoxyalkyl groups, cycloalkyl groups, alkoxy groups, haloalkoxy groups, alkylthio groups, alkylsulfonyl groups, acyl groups, haloalkylcarbonyl groups, alkoxycarbonyl groups, cyano group and carbamoyl group (its nitrogen atom may be substituted with same or different alkyl groups)].

BEST MODE FOR CARRYING OUT THE INVENTION

Although it depends on relative activities of respective ingredients, the composition of the present invention contains at least one compound of the formula (I) generally in an amount of from 0.001 to 100 parts by weight, preferably from 0.01 to 50 parts by weight, more preferably from 0.05 to 30 parts by weight to 1 part by weight of an isoxazoline derivative of the formula (I) or its salt.

One active ingredient in the composition of the present invention is a compound of the formula (I), and achieves an excellent herbicidal effect by itself.

Particularly, it does not have a substantial phytotoxicity to rice, wheat, barley, corn, grain sorghum, soybeans, cotton, sugar beet, turf, fruit trees and the like, but achieves an excellent herbicidal effect at a small dose to various weeds growing on an upland field in a wide term range of from before germination to growing season, such as gramineous weeds including barnyardgrass (*Echinochloa crusgalli* var. *crus-galli*), crabgrass (*Digitaria ciliaris*), green foxtail (*Setaria viridis*), annual bluegrass (*Poa annua*), johnsongrass (*Sorghum halepense* Pers.), blackgrass (*Alopecurus myosuroides*), wild oats (*Avena fatua*), and the like, broad leaf weeds including pale persicaria (*Polygonum lapathifolia*), slender amaranth (*Amaranthus viridis*), common lambsquarters (*Chenopodium album* L.), common chickweed (*Stellaria media* Villars), velvetleaf (*Abutilon*

theophrasti Medic), prickly sida (*Sida spinosa* L.), Hemp
sesbania (*Sesbania exaltata* Cory), common ragweed
(*Ambrosia artemisiifolia*), morningglory, and the like,
and annual and perennial sedge weeds including purple
5 nutsedge (*Cyperus rotundus* L.), yellow nutsedge (*Cyperus*
esculentus L.), hime-kugu (*Cyperus brevifolius* H.), sedge
weed (*Cyperus microiria* Steud), rice flatsedge (*Cyperus*
iria L.), and the like.

Further, the composition of the present invention
10 achieves an excellent herbicidal effect at a small dose
to various weeds growing on a paddy field in a wide term
range of from before germination to growing season, such
as annual weeds including watergrass (*Echinochloa*
oryzicola), smallflower umbrella plant (*Cyperus*
15 *difformis*), konagi (*Monochoria vaginalis*), aze-na
(*Lindernia procumbens*) and the like, and perennial weeds
including mizu-gayatsuri (*Cyperus serotinus*), kuroguwai
(*Eleocharis kuroguwai*), inu-hotaru-i (*Scirpus juncoïdes*),
and the like.

20 A compound of the following Group A which is another
active ingredient to be used in combination with an
isoxazoline derivative of the formula (I) or its salt,
has less phytotoxicity to gramineous crops such as corn
wheat and the like, but has a herbicidal activity to only
25 a part of gramineous weeds such as green foxtail and
broad leaf weeds such as barnyardgrass, common
lambsquarters and velvetleaf, and its herbicidal spectrum

is narrow.

Group A

atrazine, simazine, cyanazine, isoxaflutole,
mesotrione, flumetsulam, imazethapyr, imazapyr, dicamba,
5 clopyralid, prosulfuron, halosulfuron-methyl, rimsulfuron,
bentazone, carfentrazone-ethyl, metribuzin,
thifensulfuron-methyl, nicosulfuron, primisulfuron,
cloransulam-methyl, glufosinate, glyphosate, glyphosate-
trimesium, pendimethalin, linuron, prometryn,
10 diflufenican, flumioxazin, and metolachlor.

The present invention provides a herbicidal
composition effective for applying to a new cultivation
method such as non-tillage cultivation in order to
selectively control a wide range of weeds. The herbicidal
15 composition of the present invention is particularly
effective for killing main weeds in a corn field, such as
dicotyledons including wild buckwheat (*Fallopia*
convolvulua A.), sansevieria (*Polygonum scabrum*), common
purslane (*Portulaca oleracea* L.), common lambsquarters
20 (*Chenopodium album* L.), common amaranth (*Amaranthus*
retroflexus L.), wild mustard (*Sinapis Arvensis* L.), hemp
sesbania (*Sesbania oxaltata* Cory), sicklepod (*Cassia*
obtusifolia L.), velvetleaf (*Abutilon theophrasti* Medic),
prickly sida (*Sida spinosa* L.), ivyleaf morningglory
25 (*Ipomoea hederacea* Jacq), common morning glory (*Ipomoea*
purpurea), jimsonweed (*Datura stramonium*), black
nightshade (*Solanum nigrum* L.), common cocklebur

(*Xanthium strumarium* L.), sunflower, field bindweed
(*Convolvulus arvensis*), sun spurge (*Euphorbia*
helioscopia), devils beggarticks (*Bidens frondosa* L.),
common ragweed (*Ambrosia artemisiifolia*) and the like,
5 and monocotyledons including barnyardgrass (*Echinochloa*
crusgalli var. *crus-galli*), green foxtail (*Setaria*
viridis), giant foxtail (*Setaria faberi*), yellow foxtail
(*Setaria glauca*), crabgrass (*Digitaria ciliaris*),
goosegrass (*Eleusine indica* Gaertn), johnsongrass
10 (*Sorghum halepense* Pers.), quackgrass (*Agropyron repens*
P.), shattercane (*Sorghum vulgare*), and the like, but
does not have phytotoxicity to aimed crops such as corn
and soybeans which grow after corn.

The herbicidal composition of the present invention
15 contains at least one compound of Group A in an amount of
from 0.001 to 100 parts by weight, preferably from 0.01
to 50 parts by weight, more preferably from 0.05 to 30
parts by weight to 1 part by weight of an isoxazoline
derivative of the formula (I) or its salt. If the amount
20 of a herbicide of Group A is less than 0.001 part by
weight, a satisfactory effect can not be achieved, and if
this amount exceeds 100 parts by weight, safety to aimed
crops becomes insufficient.

Among compounds of Group A, at least one compound
25 selected from the group consisting of atrazine, cyanazine,
simazine and prometryn, or at least one compound selected
from the group consisting of glyphosate, glufosinate,

linuron and flumetsulam, is preferable. Particularly, cyanazine or atrazine is preferable.

Next, representative examples of the present compound represented by the general formula (I) are shown
5 in Tables 1 to 14. However, the present compound is not restricted to these examples.

The following abbreviated expressions used in the Tables refer to the following groups.

	Me: methyl group	Et: ethyl group
10	Pr: n-propyl group	Pr-i: isopropyl group
	Pr-c: cyclopropyl group	Bu: n-butyl group
	Bu-i: isobutyl group	Bu-s: sec-butyl group
	Bu-t: tert-butyl group	Bu-c: cyclobutyl group
	Pen: n-pentyl group	Pen-c: cyclopentyl group
15	Hex: n-hexyl group	Hex-c: cyclohexyl group
	Ph: phenyl group	

For example, (4-Cl)Ph indicates 4-chlorophenyl group, and 3-Hex indicates 3-hexyl group.

When the present compound contains hydroxyl group as
20 a substituent, there may exist keto-enol tautomers. Any of these tautomers and any mixture of these tautomers are included in the present compound.

Table 1


												
Comp. No.	R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	Z ₁	R ²²	R ²³	R ²⁴	Melting point (°C) or refractive index (n _D ²⁰)
1-0001	Me	Me	H	H	2	H	H	S	Me	H	H	66-68
1-0002	Me	Me	H	H	2	H	H	S	Cl	Me	H	87-88
1-0003	Me	Me	H	H	2	H	H	S	H	H	Me	95-97
1-0004	Me	Me	H	H	2	H	H	S	Cl	H	H	70-72
1-0005	Me	Me	H	H	2	H	H	S	H	H	Cl	118-119
1-0006	Me	Me	H	H	2	H	H	O	H	H	H	Unmesurable
1-0007	Me	Me	H	H	2	H	H	O	H	H	C(=O)OMe	124-125

Table 2


												
Comp. No.	R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	Z ²	R ²⁵	R ²⁶	R ²⁷	Melting point (°C) or refractive index (n _D ²⁰)
2-0001	Me	Me	H	H	2	H	H	S	Me	C(=NOMe)Me	Me	95-96
2-0002	Me	Me	H	H	0	H	H	S	Me	C(=NOMe)Me	Me	
2-0003	Me	Me	H	H	2	H	H	S	H	H	H	99-101
2-0004	Me	Me	H	H	2	H	H	S	H	OMe	H	96-97
2-0005	Me	Me	H	H	2	H	H	S	Cl	H	Cl	125-127
2-0006	Me	Me	H	H	2	H	H	S	Cl	Cl	Cl	158-160
2-0007	Me	Me	H	H	2	H	H	S	Me	Me	Me	117-117
2-0008	Me	Me	H	H	2	H	H	S	Me	C(=O)Me	Me	146-148
2-0009	Me	Me	H	H	2	H	H	S	Ph	C(=O)Me	Me	1.5730
2-0010	Me	Me	H	H	2	H	H	S	Ph	C(=NOMe)Me	Me	129-131
2-0011	Me	Me	H	H	2	H	H	S	Cl	C(=O)OMe	Cl	157-158
2-0012	Me	Me	H	H	2	H	H	S	Cl	C(=O)NHMe	Cl	178-180
2-0013	Me	Me	H	H	2	H	H	O	H	H	H	58-61
2-0014	Me	Me	H	H	2	H	H	O	Me	H	Cl	180-181

Table 3


											
Comp. No.	R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	R ²⁹	R ²⁸	R ³⁰	Melting point (°C) or refractive index (n _D ²⁰)
3-0001	Me	Me	H	H	0	H	H	CF ₃	Ph	Cl	89-90
3-0002	Me	Me	H	H	2	H	H	CF ₃	Ph	Cl	132-133
3-0003	Me	Me	H	H	1	H	H	Ph	Me	Cl	Unmesurable
3-0004	Me	Me	H	H	2	H	H	CF ₃	Ph	SO ₂ Et	158-160
3-0005	Me	Me	H	H	2	H	H	CF ₃	Ph	N(Me) ₂	150-151
3-0006	Me	Me	H	H	0	H	H	CF ₃	Bu-t	Cl	79-81
3-0007	Me	Me	H	H	0	H	H	CF ₃	H	Cl	120-122
3-0008	Me	Me	H	H	0	H	H	CF ₃	CHF ₂	Cl	41-42
3-0009	Me	Me	H	H	0	H	H	Cl	CHF ₂	CF ₃	89-90
3-0010	Me	Me	H	H	2	H	H	CF ₃	CHF ₂	Cl	126-127
3-0011	Me	Me	H	H	2	H	H	Cl	CHF ₂	CF ₃	136-137
3-0012	Me	Me	H	H	2	H	H	OEt	Me	CF ₃	124-125
3-0013	Me	Me	H	H	2	H	H	CF ₃	Me	OMe	113-114
3-0014	Me	Me	H	H	2	H	H	CF ₃	Me	O(2-Cl)Ph	67-70
3-0015	Me	Me	H	H	2	H	H	CF ₃	Me	OPen-c	113-114
3-0016	Me	Me	H	H	2	H	H	CF ₃	Me	CN	105-108
3-0017	Me	Me	H	H	2	H	H	Cl	Et	Cl	105-107
3-0018	Me	Me	H	H	2	H	H	CHF ₂	Me	Cl	78-79
3-0019	Me	Me	H	H	2	H	H	CF ₃	-(CH ₂) ₃ O-		151-152
3-0020	Me	Me	H	H	0	H	H	CHF ₂	Me	Cl	1.5183
3-0021	Me	Me	H	H	0	H	H	CF ₃	Ph	F	
3-0022	Me	Me	H	H	0	H	H	CF ₃	Ph	SEt	
3-0023	Me	Me	H	H	0	H	H	CF ₃	Ph	N(Me) ₂	
3-0024	Me	Me	H	H	0	H	H	OMe	Me	CF ₃	
3-0025	Me	Me	H	H	0	H	H	OH	Me	CF ₃	
3-0026	Me	Me	H	H	0	H	H	OEt	Me	CF ₃	
3-0027	Me	Me	H	H	0	H	H	CF ₃	Me	F	
3-0028	Me	Me	H	H	0	H	H	CF ₃	Me	OMe	
3-0029	Me	Me	H	H	0	H	H	CF ₃	Me	O(2-Cl)Ph	
3-0030	Me	Me	H	H	0	H	H	CF ₃	Me	OPen-c	
3-0031	Me	Me	H	H	0	H	H	CF ₃	Me	CN	
3-0032	Me	Me	H	H	0	H	H	Cl	Et	Cl	
3-0033	Me	Me	H	H	0	H	H	CF ₃	-(CH ₂) ₃ O-		
3-0034	Me	Me	H	H	2	H	H	CF ₃	H	Cl	138-140

Table 4

Comp. No.	R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	R ²⁹	R ²⁸	R ³⁰	Melting point (°C) or refractive index (n _D ²⁰)
3-0035	Me	Me	H	H	2	H	H	H	Me	Cl	105-106
3-0036	Me	Me	H	H	2	H	H	Me	Me	Me	148-150
3-0037	Me	Me	H	H	2	H	H	Me	Me	Cl	99-101
3-0038	Me	Me	H	H	2	H	H	Cl	Me	Cl	143-145
3-0039	Me	Me	H	H	2	H	H	CF ₃	Me	Cl	115-116
3-0040	Me	Me	H	H	2	H	H	Cl	Me	CF ₃	120-122
3-0041	Me	Me	H	H	2	H	H	CF ₃	Me	F	79-82
3-0042	Me	Me	H	H	2	H	H	CF ₃	Me	OH	90-92
3-0043	Me	Me	H	H	2	H	H	OMe	Me	CF ₃	125-126
3-0044	Me	Me	H	H	2	H	H	CF ₃	Me	OEt	92-94
3-0045	Me	Me	H	H	2	H	H	CF ₃	Me	OPr-i	69-71
3-0046	Me	Me	H	H	2	H	H	CF ₃	Me	OPr	82-83
3-0047	Me	Me	H	H	2	H	H	CF ₃	Me	OBu-t	86-89
3-0048	Me	Me	H	H	2	H	H	CF ₃	Me	OBu	61-62
3-0049	Me	Me	H	H	2	H	H	CF ₃	Me	OHex-c	124-125
3-0050	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ Pr-c	93-94
3-0051	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ Pen-c	112-113
3-0052	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ Hex-c	56-59
3-0053	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ C≡CH	92-93
3-0054	Me	Me	H	H	2	H	H	CF ₃	Me	OCHF ₂	129-130
3-0055	Me	Me	H	H	2	H	H	OCHF ₂	Me	CF ₃	Unmesurable
3-0056	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ CHF ₂	89-91
3-0057	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ CF ₃	93-95
3-0058	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ CN	1.4872
3-0059	Me	Me	H	H	2	H	H	CF ₃	Me	OCH ₂ Ph	79-81
3-0060	Me	Me	H	H	2	H	H	CF ₃	Me	OPh	122-123
3-0061	Me	Me	H	H	2	H	H	CF ₃	Me	O(3-Cl)Ph	Unmesurable
3-0062	Me	Me	H	H	2	H	H	CF ₃	Me	O(3-OMe)Ph	1.5059
3-0063	Me	Me	H	H	2	H	H	CF ₃	Me	O(4-Cl)Ph	68-69
3-0064	Me	Me	H	H	2	H	H	CF ₃	Me	O(4-Me)Ph	132-133
3-0065	Me	Me	H	H	2	H	H	CF ₃	Me	O(4-OMe)Ph	115-117
3-0066	Me	Me	H	H	2	H	H	CF ₃	Me	OC(=O)Me	130-131
3-0067	Me	Me	H	H	2	H	H	CF ₃	Me	SO ₂ Me	168-169
3-0068	Me	Me	H	H	2	H	H	CF ₃	Me	SEt	100-102
3-0069	Me	Me	H	H	2	H	H	CF ₃	Me	SO ₂ Et	107-108
3-0070	Me	Me	H	H	2	H	H	CF ₃	Me	SO ₂ Ph	166-168
3-0071	Me	Me	H	H	2	H	H	CF ₃	Me	Me	105-107
3-0072	Me	Me	H	H	2	H	H	Ph	Me	Cl	127-129
3-0073	Me	Me	H	H	2	H	H	CF ₃	Et	Cl	111-112

Table 5

Comp. No.	R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	R ²⁹	R ²⁸	R ³⁰	Melting point (°C) or refractive index (n _D ²⁰)
3-0074	Me	Me	H	H	2	H	H	Cl	Et	CF ₃	112-114
3-0075	Me	Me	H	H	2	H	H	CF ₃	Pr-i	Cl	157-158
3-0076	Me	Me	H	H	2	H	H	Cl	Pr-i	CF ₃	135-136
3-0077	Me	Me	H	H	2	H	H	CF ₃	Pr	Cl	89-90
3-0078	Me	Me	H	H	2	H	H	Cl	Pr	CF ₃	111-113
3-0079	Me	Me	H	H	2	H	H	CF ₃	Bu-t	H	101-103
3-0080	Me	Me	H	H	2	H	H	CF ₃	Bu-t	Cl	118-119
3-0081	Me	Me	H	H	2	H	H	CF ₃	Bu-s	Cl	110-112
3-0082	Me	Me	H	H	2	H	H	Cl	Bu-s	CF ₃	110-111
3-0083	Me	Me	H	H	2	H	H	CF ₃	Bu-i	Cl	96-98
3-0084	Me	Me	H	H	2	H	H	Cl	Bu-i	CF ₃	140-141
3-0085	Me	Me	H	H	2	H	H	CF ₃	Bu	Cl	89-90
3-0086	Me	Me	H	H	2	H	H	Cl	Bu	CF ₃	108-110
3-0087	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Ph	Cl	132-133
3-0088	Me	Me	H	H	2	H	H	Cl	CH ₂ Ph	CF ₃	118-120
3-0089	Me	Me	H	H	2	H	H	CF ₃	Pen-c	Cl	130-131
3-0090	Me	Me	H	H	2	H	H	Cl	Pen-c	CF ₃	147-148
3-0091	Me	Me	H	H	2	H	H	CF ₃	Hex-c	Cl	151-152
3-0092	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Pr-c	Cl	93-95
3-0093	Me	Me	H	H	2	H	H	Cl	CH ₂ Pr-c	CF ₃	129-130
3-0094	Me	Me	H	H	2	H	H	CF ₃	1-cyclopropylethyl	Cl	87-89
3-0095	Me	Me	H	H	2	H	H	Cl	1-cyclopropylethyl	CF ₃	121-123
3-0096	Me	Me	H	H	2	H	H	CF ₃	CH ₂ (2-Methylcyclopropyl)	Cl	102-103
3-0097	Me	Me	H	H	2	H	H	Cl	CH ₂ (2-Methylcyclopropyl)	CF ₃	118-119
3-0098	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Bu-c	Cl	94-96
3-0099	Me	Me	H	H	2	H	H	Cl	CH ₂ Bu-c	CF ₃	141-142
3-0100	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Pen-c	Cl	127-129
3-0101	Me	Me	H	H	2	H	H	Cl	CH ₂ Pen-c	CF ₃	146-149
3-0102	Me	Me	H	H	2	H	H	CF ₃	CH ₂ Hex-c	Cl	152-154
3-0103	Me	Me	H	H	2	H	H	Cl	CH ₂ Hex-c	CF ₃	115-117
3-0104	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CH=CH ₂	Cl	78-80
3-0105	Me	Me	H	H	2	H	H	Cl	CH ₂ CH=CH ₂	CF ₃	105-106
3-0106	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C≡CH	Cl	73-74
3-0107	Me	Me	H	H	2	H	H	Cl	CH ₂ C≡CH	CF ₃	108-109
3-0108	Me	Me	H	H	2	H	H	CF ₃	CHMeC≡CH	Cl	95-96
3-0109	Me	Me	H	H	2	H	H	Cl	CHMeC≡CH	CF ₃	116-118
3-0110	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C≡CMe	Cl	114-115
3-0111	Me	Me	H	H	2	H	H	Cl	CH ₂ C≡CMe	CF ₃	115-116
3-0112	Me	Me	H	H	2	H	H	CF ₃	CHF ₂	OMe	72-74

Table 6

Comp. No.	R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	R ²⁹	R ²⁸	R ³⁰	Melting point (°C) or refractive index (n _D ²⁰)
3-0113	Me	Me	H	H	2	H	H	OMe	CHF ₂	CF ₃	108-109
3-0114	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CHF ₂	Cl	99-100
3-0115	Me	Me	H	H	2	H	H	Cl	CH ₂ CHF ₂	CF ₃	107-109
3-0116	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CF ₃	Cl	135-136
3-0117	Me	Me	H	H	2	H	H	Cl	CH ₂ CF ₃	CF ₃	112-115
3-0118	Me	Me	H	H	2	H	H	CF ₃	CH ₂ OMe	Cl	87-89
3-0119	Me	Me	H	H	2	H	H	Cl	CH ₂ OMe	CF ₃	125-128
3-0120	Me	Me	H	H	2	H	H	CF ₃	CH ₂ OEt	Cl	97-98
3-0121	Me	Me	H	H	2	H	H	Cl	CH ₂ OEt	CF ₃	128-129
3-0122	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CH ₂ OH	Cl	79-81
3-0123	Me	Me	H	H	2	H	H	Cl	CH ₂ CH ₂ OH	CF ₃	93-94
3-0124	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CH ₂ OMe	Cl	102-104
3-0125	Me	Me	H	H	2	H	H	Cl	CH ₂ CH ₂ OMe	CF ₃	118-119
3-0126	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CH ₂ OEt	Cl	56-59
3-0127	Me	Me	H	H	2	H	H	Cl	CH ₂ CH ₂ OEt	CF ₃	118-119
3-0128	Me	Me	H	H	2	H	H	CF ₃	CH ₂ SMe	Cl	103-105
3-0129	Me	Me	H	H	2	H	H	Cl	CH ₂ SMe	CF ₃	128-129
3-0130	Me	Me	H	H	2	H	H	CF ₃	CH ₂ SO ₂ Me	Cl	157-159
3-0131	Me	Me	H	H	2	H	H	Cl	CH ₂ SO ₂ Me	CF ₃	165-166
3-0132	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CH ₂ SO ₂ Me	Cl	155-157
3-0133	Me	Me	H	H	2	H	H	Cl	CH ₂ CH ₂ SO ₂ Me	CF ₃	166-168
3-0134	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CN	Cl	128-129
3-0135	Me	Me	H	H	2	H	H	Cl	CH ₂ CN	CF ₃	117-118
3-0136	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C(=O)OEt	Cl	127-129
3-0137	Me	Me	H	H	2	H	H	Cl	CH ₂ C(=O)OEt	CF ₃	143-145
3-0138	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C(=O)NH ₂	Cl	173-174
3-0139	Me	Me	H	H	2	H	H	Cl	CH ₂ C(=O)NH ₂	CF ₃	182-183
3-0140	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C(=O)N(Me) ₂	Cl	142-143
3-0141	Me	Me	H	H	2	H	H	Cl	CH ₂ C(=O)N(Me) ₂	CF ₃	181-182
3-0142	Me	Me	H	H	2	H	H	CF ₃	CH ₂ C(=O)Me	Cl	148-149
3-0143	Me	Me	H	H	2	H	H	Cl	CH ₂ C(=O)Me	CF ₃	163-164
3-0144	Me	Me	H	H	2	H	H	CF ₃	CH ₂ CH ₂ C(=O)Me	Cl	89-91
3-0145	Me	Me	H	H	2	H	H	Me	Ph	Me	140-141
3-0146	Me	Me	H	H	2	H	H	Me	Ph	Cl	124-125
3-0147	Me	Me	H	H	2	H	H	Et	Ph	Cl	112-113
3-0148	Me	Me	H	H	2	H	H	Pr	Ph	Cl	122-123
3-0149	Me	Me	H	H	2	H	H	Pr-i	Ph	Cl	116-117
3-0150	Me	Me	H	H	2	H	H	Bu-t	Ph	Cl	100-102
3-0151	Me	Me	H	H	2	H	H	CF ₃	Ph	H	111-112

Table 7

Comp. No.	R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	R ²⁹	R ²⁸	R ³⁰	Melting point (°C) or refractive index (n _D ²⁰)
3-0152	Me	Me	H	H	2	H	H	CF ₃	Ph	Me	129-132
3-0153	Me	Me	H	H	2	H	H	CF ₃	Ph	CF ₃	112-113
3-0154	Me	Me	H	H	2	H	H	CF ₃	Ph	F	90-91
3-0155	Me	Me	H	H	2	H	H	CF ₃	Ph	OMe	104-106
3-0156	Me	Me	H	H	2	H	H	CF ₃	Ph	OEt	129-131
3-0157	Me	Me	H	H	2	H	H	CF ₃	Ph	OPr-i	86-88
3-0158	Me	Me	H	H	2	H	H	CF ₃	Ph	OPr	117-118
3-0159	Me	Me	H	H	2	H	H	CF ₃	Ph	OBu-t	105-108
3-0160	Me	Me	H	H	2	H	H	CF ₃	Ph	OCHF ₂	90-92
3-0161	Me	Me	H	H	2	H	H	CF ₃	Ph	SO ₂ Me	167-168
3-0162	Me	Me	H	H	2	H	H	CF ₃	Ph	CN	113-115
3-0163	Me	Me	H	H	2	H	H	CF ₃	(2-Cl)Ph	Cl	153-154
3-0164	Me	Me	H	H	2	H	H	CF ₃	(3-Cl)Ph	Cl	106-107
3-0165	Me	Me	H	H	2	H	H	CF ₃	(4-Cl)Ph	Cl	142-143
3-0166	Me	Me	H	H	2	H	H	CF ₃	(4-F)Ph	Cl	135-138
3-0167	Me	Me	H	H	2	H	H	CF ₃	(4-OMe)Ph	Cl	136-138
3-0168	Me	Me	H	H	2	H	H	CF ₃	(4-Me)Ph	Cl	129-130
3-0169	Me	Me	H	H	2	H	H	CF ₃	(4-NO ₂)Ph	Cl	145-147
3-0170	Me	Me	H	H	2	H	H	CF ₃	(4-CN)Ph	Cl	91-93
3-0171	Me	Me	H	H	2	H	H	CF ₃	(4-C(=O)Me)Ph	Cl	133-135
3-0172	Me	Me	H	H	2	H	H	CF ₃	(4-C(=O)OMe)Ph	Cl	121-124
3-0173	Me	Me	H	H	2	H	H	CF ₃	Pyrimidin-2-yl	Cl	148-150
3-0174	Me	Me	H	H	2	H	H	CF ₃	4,6-Dimethoxy-pyrimidin-2-yl	Cl	117-118
3-0175	Me	Me	H	H	2	H	H	CF ₃	SO ₂ Me	Cl	146-148
3-0176	Me	Me	H	H	2	H	H	CF ₃	SO ₂ Ph	Cl	145-148
3-0177	Me	Me	H	H	2	H	H	CF ₃	C(=O)Me	Cl	130-131
3-0178	Me	Me	H	H	2	H	H	CF ₃	C(=O)Ph	Cl	114-117
3-0179	Me	Me	H	H	2	H	H	CF ₃	C(=O)OMe	Cl	104-106
3-0180	Me	Et	H	H	2	H	H	CF ₃	Me	Cl	108-110
3-0181	Me	Me	H	H	0	H	H	CHF ₂	Me	Cl	1.5183
3-0182	Me	Me	H	H	0	H	H	Ph	Me	Cl	76-77
3-0183	Me	Me	H	H	0	H	H	CF ₃	Bu-t	OMe	1.4831
3-0184	Me	Me	H	H	0	H	H	CF ₃	CH ₂ C(=O)NH ₂	Cl	179-180
3-0185	Me	Me	H	H	0	H	H	Me	Ph	Cl	58-60
3-0186	Me	Me	H	H	0	H	H	CF ₃	Me	Cl	
3-0187	Me	Me	H	H	0	H	H	CF ₃	Me	OCHF ₂	
3-0188	Me	Me	H	H	2	H	H	CF ₃	Me	OCHF ₂	129-130
3-0189	Me	Me	H	H	0	H	H	CF ₃	Et	OCHF ₂	
3-0190	Me	Me	H	H	2	H	H	CF ₃	Et	OCHF ₂	98-100

Table 8


											
Comp. No.	R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	Z ³	R ³¹	R ³²	Melting point (°C) or refractive index (n _D ²⁰)
4-0001	Me	Me	H	H	2	H	H	O	CF ₃	Me	135-136
4-0002	Me	Me	H	H	2	H	H	S	Me	Cl	113-114
4-0003	Me	Me	H	H	0	H	H	O	CF ₃	Me	
4-0004	Me	Me	H	H	0	H	H	S	Me	Cl	
4-0005	Me	Me	H	H	2	H	H	O	Me	Me	178-179
4-0006	Me	Me	H	H	2	H	H	O	CF ₃	OEt	89-91
4-0007	Me	Me	H	H	2	H	H	O	Ph	Me	81-83
4-0008	Me	Me	H	H	2	H	H	S	Me	OEt	109-111

Table 9


											
Comp. No.	R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	Z ⁴	R ³³	R ³⁴	Melting point (°C) or refractive index (n _D ²⁰)
5-0001	Me	Me	H	H	2	H	H	NMe	Cl	Me	114-115
5-0002	Me	Me	H	H	2	H	H	NMe	Cl	Et	107-108
5-0003	Me	Me	H	H	2	H	H	NMe	CF ₃	H	142-143
5-0004	Me	Me	H	H	2	H	H	NCHF ₂	-(CH ₂) ₄ -		123-125
5-0005	Me	Me	H	H	2	H	H	NPh	OEt	Me	1.5397
5-0006	Me	Me	H	H	2	H	H	NPh	OCHF ₂	Me	1.5339
5-0007	Me	Me	H	H	2	H	H	NPh	CF ₃	H	99-101
5-0008	Me	Me	H	H	2	H	H	NPh	OCH ₂ CH=CH ₂	Me	87-90
5-0009	Me	Me	H	H	1	H	H	NPh	OCH ₂ CH=CH ₂	Me	1.5702

Table 10


											
Comp. No.	R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	Z ⁵	R ³⁵	R ³⁶	Melting point (°C) or refractive index (n _D ²⁰)
6-0001	Me	Me	H	H	2	H	H	NCHF ₂	-(CH ₂) ₄ -		Unmesurable
6-0002	Me	Me	H	H	2	H	H	NPh	H	OEt	107-108
6-0003	Me	Me	H	H	2	H	H	NPh	H	OCHF ₂	1.5383
6-0004	Me	Me	H	H	2	H	H	O	Me	H	100-102
6-0005	Me	Me	H	H	0	H	H	NCHF ₂	-(CH ₂) ₄ -		1.5264

Table 11


													
Comp. No.	R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	R ³⁷	R ³⁸	R ³⁹	R ⁴⁰		Melting point (°C) or refractive index (n _D ²⁰)
7-0001	Me	Me	H	H	2	H	H	H	CF ₃	H	H	-	77-80
7-0002	Me	Me	H	H	2	H	H	H	CF ₃	H	H	N-oxide	114-116
7-0003	Me	Me	H	H	0	H	H	H	CF ₃	H	H	-	
7-0004	Me	Me	H	H	2	H	H	H	H	H	H	-	130-131
7-0005	Me	Me	H	H	2	H	H	H	H	H	H	N-oxide	166-168
7-0006	Me	Me	H	H	2	H	H	Cl	Ph	H	H	-	118-120
7-0007	Me	Me	H	H	2	H	H	OMe	Ph	H	H	-	105-106
7-0008	Me	Me	H	H	2	H	H	Cl	Me	H	H	-	115-116
7-0009	Me	Me	H	H	2	H	H	OMe	Me	H	H	-	134-135
7-0010	Me	Me	H	H	2	H	H	Me	Me	H	H	N-oxide	198-199
7-0011	Me	Me	H	H	2	H	H	Ph	Ph	H	H	-	161-162
7-0012	Me	Me	H	H	1	H	H	H	H	H	H	-	97-99
7-0013	Me	Me	H	H	0	H	H	(2-Chloropyridin-3-yl)methylthio	H	H	H	-	154-155

Table 12


											
Comp. No.	R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	R ⁴¹	R ⁴²	R ⁴³	Melting point (°C) or refractive index (n _D ²⁰)
8-0001	Me	Me	H	H	2	H	H	H	OMe	CF ₃	175-176
8-0002	Me	Me	H	H	0	H	H	H	OMe	CF ₃	
8-0003	Me	Me	H	H	2	H	H	H	Cl	Cl	119-120
8-0004	Me	Me	H	H	2	H	H	H	OEt	CF ₃	94-95
8-0005	Me	Me	H	H	2	H	H	H	OMe	OMe	186-187
8-0006	Me	Me	H	H	2	H	H	Me	OMe	CF ₃	143-144
8-0007	Me	Me	H	H	2	H	H	OMe	OMe	CF ₃	144-145
8-0008	Me	Me	H	H	2	H	H	SMe	OMe	CF ₃	160-162
8-0009	Me	Me	H	H	2	H	H	SO ₂ Me	OMe	CF ₃	144-146
8-0010	Me	Me	H	H	2	H	H	NH ₂	OMe	CF ₃	208-209
8-0011	Me	Me	H	H	2	Pr-i	H	H	H	CF ₃	112-113
8-0012	Me	Me	H	H	0	Pr-i	H	H	H	CF ₃	1.4986

Table 13



									
Comp. No.	R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	Y ¹	Melting point (°C) or refractive index (n _D ²⁰)
9-0001	Me	Me	H	H	2	H	H	Pyridin-2-yl	116-118
9-0002	Me	Me	H	H	2	H	H	Pyridin-2-yl 1-oxide	140-143
9-0003	Me	Me	H	H	2	H	H	Pyridin-4-yl	133-136
9-0004	Me	Me	H	H	2	H	H	Pyridin-4-yl 1-oxide	110-113
9-0005	Me	Me	H	H	2	H	H	1,2,4-Oxadiazol-3-yl	Unmesurable
9-0006	Me	Me	H	H	2	H	H	3-Phenyl-1,2,4-oxadiazol-5-yl	153-154
9-0007	Me	Me	H	H	2	H	H	3-Benzyl-1,2,4-oxadiazol-5-yl	108-109
9-0008	Me	Me	H	H	2	H	H	2-Chlorothiazol-4-yl	110-112
9-0009	Me	Me	H	H	2	H	H	1,4-Dimethylimidazol-5-yl	163-164
9-0010	Me	Me	H	H	1	H	H	Pyridin-2-yl	81-82
9-0011	Me	Me	H	H	1	H	H	Pyridin-4-yl	94-96
9-0012	Me	Me	H	H	1	H	H	1,4-Dimethylimidazol-5-yl	138-140
9-0013	Me	Me	H	H	0	H	H	1,4-Dimethylimidazol-5-yl	1.5427

Table 14

									
Comp. No.	R ¹	R ²	R ³	R ⁴	n	R ⁵	R ⁶	Y ¹	Melting point (°C) or refractive index (n _D ²⁰)
10-0001	Me	Me	H	H	2	H	H	Benzimidazol-2-yl	171-174
10-0002	Me	Me	H	H	2	H	H	Benzothiophen-2-yl	181-183
10-0003	Me	Me	H	H	2	H	H	3-Chlorobenzothiophen-2-yl	109-112
10-0004	Me	Me	H	H	2	H	H	Benzotriazol-1-yl	206-207
10-0005	Me	Me	H	H	2	H	H	1-Methylindazol-4-yl	128-130
10-0006	Me	Me	H	H	2	H	H	Benzothiazol-2-yl	142-143
10-0007	Me	Me	H	H	2	H	H	Benzothiophen-3-yl	188-191
10-0008	Me	Me	H	H	2	H	H	5-Chlorobenzothiophen-3-yl	129-130
10-0009	Me	Me	H	H	2	H	H	Benoxazol-2-yl	127-129
10-0010	Me	Me	H	H	2	H	H	3-Methylbenzothiophen-2-yl	161-163
10-0011	Me	Me	H	H	2	H	H	3-Bromobenzothiophen-2-yl	118-119
10-0012	Me	Me	H	H	2	H	H	Benzofuran-2-yl	123-124
10-0013	Me	Me	H	H	2	H	H	2-Methylbenzofuran-7-yl	135-137
10-0014	Me	Me	H	H	2	H	H	3-Bromobenzofuran-2-yl	107-108
10-0015	Me	Me	H	H	2	H	H	Benzothiophen-7-yl	95-97
10-0016	Me	Me	H	H	2	H	H	1-Methylindazol-7-yl	89-90
10-0017	Me	Me	H	H	2	H	H	3-Methylbenzofuran-2-yl	111-112
10-0018	Me	Me	H	H	2	H	H	3-Chloro-1-methylindol-2-yl	162-165

In using the composition of the present invention as a herbicide, the present composition may be used in a mixture by itself without adding other compounds, but it can also be used in the form of a wettable powder, granules, fine granules, a powder, an emulsifiable concentrate, a solution, a suspension, a flowable, etc. by mixing with a carrier, a surfactant, a dispersant, an adjuvant, etc. all generally usable in formulation.

As the carrier usable in formulation, there can be mentioned, for example, solid carriers such as talc, bentonite, clay, kaolin, diatomaceous earth, white carbon, vermiculite, calcium carbonate, slaked lime, siliceous sand, ammonium sulfate, urea and the like; and liquid carriers such as isopropyl alcohol, xylene, cyclohexane, methylnaphthalene and the like.

As the surfactant and the dispersant, there can be mentioned, for example, metal salts of alkylbenzenesulfonic acids, metal salts of an alkylnaphthalenesulfonic acid-formalin condensate, salts of alcohol sulfates, alkylarylsulfonic acid salts, ligninsulfonic acid salts, polyoxyethylene glycol ethers, polyoxyethylene alkyl aryl ethers, monoalkylates of polyoxyethylene sorbitan and the like. As the adjuvant, there can be mentioned, for example, carboxymethyl cellulose, polyethylene glycol and gum arabic.

Also, the composition of the present invention may be prepared by formulating respective active ingredients

in the above-mentioned manner and then mixing the respective ingredients. The composition of the present invention thus formulated may be applied to plants as it is or may be applied after diluting with water or the like. The composition of the present invention may be mixed with other herbicides to enhance its herbicidal effect, or may be used further in combination with an insecticide, a fungicide, a plant growth-regulating agent, a fertilizer, a soil-improving agent or the like.

The composition of the present invention contains an isoxazoline derivative of the formula (I) or its salt and at least one compound selected from Group A preferably in a total amount of from 0.5 to 90 wt%, more preferably from 1 to 80 wt%.

Also, an isoxazoline derivative of the formula (I) or its salt may be mixed with two or more compounds selected from Group A, and its application amount is preferably from 0.5 to 90 wt%, more preferably from 1 to 80 wt%, in a total amount of the two ingredients.

The compound of the formula (I) used in the composition of the present invention can be prepared as illustrated in the following Preparation Examples, but should not be limited thereto.

<Preparation Example 1>

Production of 3-(5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0001)

2.1 g of sodium hydrosulfide hydrate (purity: 70%, 26.2 mmol) was added to a solution of 2.3 g (13.1 mmol) of 3-methylsulfonyl-5,5-dimethyl-2-isoxazoline dissolved in 20 ml of N,N-dimethylformamide. The mixture was stirred for 2 hours. Thereafter were added 1.8 g (13.1 mmol) of anhydrous potassium carbonate, 2.0 g (13.1 mmol) of Rongalit and 3.6 g (10.5 mmol) of 4-bromomethyl-5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazole. The resulting mixture was stirred at room temperature for 15 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 2.7 g (yield: 65.5%) of 3-(5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline as white crystals (melting point: 89 to 90°C).

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

7.55-7.50 (5H,m), 4.33 (2H,s), 2.83 (2H,s),
1.45 (6H,s)

<Preparation Example 2>

Production of 3-(5-chloro-1-phenyl-3-trifluoromethyl-1H-

pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline
(present compound No. 3-0002)

0.63 g of m-chloroperbenzoic acid (purity: 70%, 2.6
mmoles) was added, with ice-cooling, to a solution of 0.4
5 g (1.0 mmoles) of 3-(5-chloro-1-phenyl-3-trifluoromethyl-
1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline
dissolved in 15 ml of chloroform. The mixture was
stirred at room temperature for 22 hours to give rise to
a reaction. After the completion of the reaction, the
10 reaction mixture was poured into water, followed by
extraction with chloroform. The resulting organic layer
was washed with an aqueous sodium hydrogensulfite
solution, an aqueous sodium hydrogencarbonate solution
and an aqueous sodium chloride solution in this order and
15 then dried over anhydrous magnesium sulfate. The
resulting solution was subjected to vacuum distillation
to remove the solvent contained therein. The resulting
crystals were washed with hexane to obtain 0.4 g (yield:
83.2%) of 3-(5-chloro-1-phenyl-3-trifluoromethyl-1H-
20 pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline as
white crystals (melting point: 132 to 133°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.60-7.51 (5H,m), 4.37 (2H,s), 3.14 (2H,s) 1.53
(6H,s)

25 <Preparation Example 3>

Production of 3-(5-chloro-1-methyl-3-phenyl-1H-pyrazol-4-
ylmethylsulfinyl)-5,5-dimethyl-2-isoxazoline (present

compound No. 3-0003)

0.87 g of m-chloroperbenzoic acid (purity: 70%, 3.54 mmols) was added, with ice-cooling, to a solution of 0.85 g (2.53 mmols) of 3-(5-chloro-1-methyl-3-phenyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline dissolved in 30 ml of chloroform. The mixture was stirred at room temperature for 1 hour to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 0.48 g (yield: 53.9%) of 3-(5-chloro-1-methyl-3-phenyl-1H-pyrazol-4-ylmethylsulfinyl)-5,5-dimethyl-2-isoxazoline as a transparent viscous substance.

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

7.63-7.60 (2H,m), 7.48-7.37 (3H,m), 4.29 (2H,q),
3.91 (3H,s), 3.12 (1H,d), 2.79 (1H,d), 1.41 (3H,s),
1.35 (3H,s)

<Preparation Example 4>

Production of 5,5-dimethyl-3-(5-fluoro-1-phenyl-3-

trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline
(present compound No. 3-0021)

9.3 g of sodium hydrosulfide hydrate (purity: 70%,
116.3 mmol) was added to a solution of 18.7 g (105.7
5 mmol) of 5,5-dimethyl-3-methylsulfonyl-2-isoxazoline
(present compound No. 2-1) dissolved in 300 ml of N,N-
dimethylformamide. The mixture was stirred for 2 hours.
The reaction system was ice-cooled. Thereto was added a
solution of 30.3 g (93.8 mmol) of 4-bromomethyl-5-
10 fluoro-1-phenyl-3-trifluoromethyl-1H-pyrazole dissolved
in 200 ml of N,N-dimethylformamide. The mixture was
stirred at 0°C for 30 minutes to give rise to a reaction.
After the completion of the reaction, the reaction
mixture was poured into water, followed by extraction
15 with ethyl acetate. The resulting organic layer was
washed with an aqueous sodium chloride solution and then
dried over anhydrous magnesium sulfate. The resulting
solution was subjected to vacuum distillation to remove
the solvent contained therein. The residue was purified
20 by silica gel column chromatography (developing solvent:
hexane-ethyl acetate mixed solvent) to obtain 13.11 g
(yield: 37.4%) of 5,5-dimethyl-3-(5-fluoro-1-phenyl-3-
trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline
as a yellow oily substance.

25 ¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.65-7.39 (5H,m), 4.24 (2H,s), 2.81 (2H,s),
1.43 (6H,s)

<Preparation Example 5>

Production of 5,5-dimethyl-3-(5-ethylthio-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline (present compound No. 3-0022)

5 0.2 g (4.0 mmoles) of sodium hydroxide and 1 ml of water were added to a solution of 0.25 g (4.0 mmoles) of ethanethiol dissolved in 10 ml of N,N-dimethylformamide. The mixture was stirred at room temperature for 30 minutes. Thereto was added a solution of 0.5 g (1.4
10 mmoles) of 5,5-dimethyl-3-(5-fluoro-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline dissolved in 5 ml of N,N-dimethylformamide. The resulting mixture was stirred for 1 hour to give rise to a reaction. After the completion of the reaction, the
15 reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation
20 to remove the solvent contained therein, to obtain 0.6 g (yield: 100%) of 5,5-dimethyl-3-(5-ethylthio-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.62-7.47 (5H,m), 4.44 (2H,s), 2.83 (2H,s),
25 2.50 (2H,q), 1.45 (6H,s), 1.02 (3H,t)

<Preparation Example 6>

Production of 5,5-dimethyl-3-(5-ethylsulfonyl-1-phenyl-3-

trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-2-isoxazoline (present compound No. 3-0004)

1.7 g of m-chloroperbenzoic acid (purity: 70%, 6.7 mmoles) was added, with ice-cooling, to a solution of 0.6 g (1.3 mmoles) of 5,5-dimethyl-3-(5-ethylthio-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline dissolved in 10 ml of chloroform. The mixture was stirred at room temperature for 16 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The resulting crystals were washed with hexane to obtain 0.6 g (yield: 93.0%) of 5,5-dimethyl-3-(5-ethylsulfonyl-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-2-isoxazoline as light yellow crystals (melting point: 158 to 160°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.58-7.54 (5H,m), 5.16 (2H,s), 3.18 (2H,s),
3.15 (2H,q), 1.55 (6H,s), 1.24 (3H,t)

<Preparation Example 7>

Production of 5,5-dimethyl-3-(5-dimethylamino-1-phenyl-3-

trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline
(present compound 3-0023)

0.8 g (6.7 mmoles) of a 40% aqueous dimethylamine
solution was added to a solution of 0.5 g (1.3 mmoles) of
5 5,5-dimethyl-3-(5-fluoro-1-phenyl-3-trifluoromethyl-1H-
pyrazol-4-ylmethylthio)-2-isoxazoline dissolved in 10 ml
of N,N-dimethylformamide. The mixture was stirred at
100°C for 9 hours in a sealed tube. Thereto was added
3.0 g (26.6 mmoles) of a 40% aqueous dimethylamine
10 solution, and the resulting mixture was stirred for 9
hours to give rise to a reaction. After the completion
of the reaction, the reaction mixture was poured into
water, followed by extraction with ethyl acetate. The
resulting organic layer was washed with an aqueous sodium
15 chloride solution and then dried over anhydrous sodium
sulfate. The resulting solution was subjected to vacuum
distillation to remove the solvent contained therein.
The residue was purified by silica gel column
chromatography (developing solvent: hexane-ethyl acetate
20 mixed solvent) to obtain 0.4 g (yield: 80.6%) of 5,5-
dimethyl-3-(5-dimethylamino-1-phenyl-3-trifluoromethyl-
1H-pyrazol-4-ylmethylthio)-2-isoxazoline.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.58-7.38 (5H,m), 4.35 (2H,s), 2.82 (2H,s),
25 2.77 (6H,s), 1.45 (6H,s)

<Preparation Example 8>

Production of 5,5-dimethyl-3-(5-dimethylamino-1-phenyl-3-

trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-2-isoxazoline (present compound 3-0005)

0.7 g of m-chloroperbenzoic acid (purity: 70%, 2.7 mmols) was added, with ice-cooling, to a solution of 0.4 g (1.1 mmols) of 5,5-dimethyl-3-(5-dimethylamino-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline dissolved in 10 ml of chloroform. The mixture was stirred at room temperature for 20 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The resulting crystals were washed with hexane to obtain 0.2 g (yield: 52.0%) of 5,5-dimethyl-3-(5-dimethylamino-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-2-isoxazoline as a white powder (melting point: 150 to 151°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.61-7.38 (5H,m), 4.75 (2H,s), 3.13 (2H,s),
2.76 (6H,s), 1.53 (6H,s)

<Preparation Example 9>

Production of 3-(1-tert-butyl-5-chloro-3-trifluoromethyl-

1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline
(present compound No. 3-0006)

21.8 g of sodium hydrosulfide (purity: 70%, 272.5
mmoles) was added to a solution of 24.1 g (136.0 mmoles)
5 of 5,5-dimethyl-3-methylsulfonyl-2-isoxazoline dissolved
in 200 ml of N,N-dimethylformamide. The mixture was
stirred for 1 hour. Thereto were added 18.8 g (136.2
mmoles) of anhydrous potassium carbonate and 21.0 g
(136.2 mmoles) of Rongalit. The resulting mixture was
10 stirred for 2 hours. Thereto was added, with ice-cooling,
40 g (125 mmoles) of 4-bromomethyl-1-tert-butyl-5-chloro-
3-trifluoromethyl-1H-pyrazole. The resulting mixture was
stirred at room temperature for 2 hours to give rise to a
reaction. After the completion of the reaction, the
15 reaction mixture was poured into water, followed by
extraction with ethyl acetate. The resulting organic
layer was washed with water and an aqueous sodium
chloride solution and then dried over anhydrous magnesium
sulfate. The resulting solution was subjected to vacuum
20 distillation to remove the solvent contained therein.
The residue was purified by silica gel column
chromatography (developing solvent: hexane-ethyl acetate
mixed solvent) to obtain 23.0 g (yield: 57.1%) of 3-(1-
tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazol-4-
25 ylmethylthio)-5,5-dimethyl-2-isoxazoline as light pink
crystals (melting point: 79.0 to 81.0°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.24 (2H,s), 2.80 (2H,s), 1.71 (9H,s), 1.43 (6H,s)

<Preparation Example 10>

Production of 3-(5-chloro-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline (present

5 compound No. 3-0007)

19.8 g (53.4 mmoles) of 3-(1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline was added to 170 ml of a 25% hydrogen bromide-acetic acid solution. The mixture was stirred at
10 40 to 50°C for 2 hours to give rise to a reaction. After the completion of the reaction was confirmed, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium
15 chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 12.0 g (yield: 60.6%) of 3-(5-chloro-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-
20 2-isoxazoline as light yellow crystals (melting point: 120.0 to 122.0°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.26 (2H,s), 2.81 (2H,s), 1.44 (6H,s)

<Preparation Example 11>

25 Production of 3-(5-chloro-1-difluoromethyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0008) and 3-(3-

chloro-1-difluoromethyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline (present compound 3-0009)

3.1 g (22.5 mmoles) of anhydrous potassium carbonate
5 was added to a solution of 2.3 g (7.3 mmoles) of 3-(5-chloro-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline dissolved in 50 ml of N,N-dimethylformamide. Thereinto was blown
chlorodifluoromethane. The resulting mixture was stirred
10 at 130 to 140°C for 3 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was pored into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium
15 chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate
20 mixed solvent) to obtain 0.69 g (yield: 25.8%) of 3-(5-chloro-1-difluoromethyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline as light yellow crystals (melting point: 41.0 to 42.0°C) and 0.54 g (yield: 20.2%) of 3-(3-chloro-1-difluoromethyl-5-
25 trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline as a white powder (melting point: 89.0 to 90.0°C).

3-(5-Chloro-1-difluoromethyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.22 (1H,t), 4.25 (2H,s), 2.80 (2H,s), 1.44 (6H,s)

5 3-(3-Chloro-1-difluoromethyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.19 (1H,t), 4.28 (2H,s), 2.80 (2H,s), 1.44 (6H,s)

<Preparation Example 12>

10 Production of 3-(5-chloro-1-difluoromethyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0010)

1.4 g of m-chloroperbenzoic acid (purity: 70%, 8.1 mmoles) was added, with ice-cooling, to a solution of
15 0.69 g (1.9 mmoles) of 3-(5-chloro-1-difluoromethyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline dissolved in 20 ml of chloroform. The mixture was stirred for 1 hour and then at room temperature for 12 hours to give rise to a reaction.
20 After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution,
25 water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum

distillation to remove the solvent contained therein.

The resulting solid was washed with n-hexane to obtain 0.4 g (yield: 53.3%) of 3-(5-chloro-1-difluoromethyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline as a white powder (melting point: 126.0 to 127.0°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.26 (1H,t), 4.68 (2H,s), 3.11 (2H,s), 1.53 (6H,s)

<Preparation Example 13>

10 Production of 3-(3-chloro-1-difluoromethyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0011)

1.1 g of m-chloroperbenzoic acid (purity: 70%, 6.4 mmoles) was added, with ice-cooling, to a solution of 0.54 g (1.5 mmoles) of 3-(3-chloro-1-difluoromethyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline dissolved in 20 ml of chloroform. The mixture was stirred for 1 hour and then at room temperature for 12 hours to give rise to a reaction.

20 After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution, 25 water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum

distillation to remove the solvent contained therein.

The resulting solid was washed with n-hexane to obtain 0.47 g (yield: 79.7%) of 3-(3-chloro-1-difluoromethyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline as a white powder (melting point: 136.0 to 137.0°C).

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

7.23 (1H,t), 4.71 (2H,s), 3.11 (2H,s), 1.53 (6H,s)

<Preparation Example 14>

10 Production of 5,5-dimethyl-3-(3-methoxy-1-methyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline (present compound No. 3-0024)

3.1 g of sodium hydrosulfide hydrate (purity: 70%, 22.0 mmoles) was added to a solution of 3.3 g (17.3 mmoles) of 5,5-dimethyl-3-ethylsulfonyl-2-isoxazoline dissolved in 10 ml of N,N-dimethylformamide. The mixture was stirred for 2 hours. Thereto were added 3.1 g (22.0 mmoles) of anhydrous potassium carbonate, 2.7 g (17.5 mmoles) of Rongalit and 4.0 g (17.5 mmoles) of 4-chloromethyl-3-methoxy-1-methyl-5-trifluoromethyl-1H-pyrazole. The resulting mixture was stirred at room temperature for 2 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting

20

25

solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 2.8 g

5 (yield: 52.0%) of 5,5-dimethyl-3-(3-methoxy-1-methyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline.
<Preparation Example 15>

Production of 5,5-dimethyl-3-(3-hydroxy-1-methyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline
10 (present compound No. 3-0025)

To 20 ml of a 25% hydrogen bromide acetic acid solution was added 3.3 g (10.6 mmoles) of 5,5-dimethyl-3-(3-methoxy-1-methyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline. The mixture was stirred at
15 50°C for 3 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was subjected to vacuum distillation to remove the solvent contained therein. The residue was poured into water. The resulting crystals were collected by filtration,
20 washed with water and dried to obtain 3.1 g (yield: 96.0%) of intended 5,5-dimethyl-3-(3-hydroxy-1-methyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline.
<Preparation Example 16>

Production of 5,5-dimethyl-3-(3-ethoxy-1-methyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline
25 (present compound No. 3-0026)

0.20 g (1.3 mmoles) of anhydrous potassium carbonate

and 0.20 g (1.5 mmoles) of ethyl iodide were added to a solution of 0.30 g (1.0 mmoles) of 5,5-dimethyl-3-(3-hydroxy-1-methyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline dissolved in 10 ml of N,N-dimethylformamide. The mixture was stirred at 50°C for 3 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 0.30 g (yield: 92.0%) of intended 5,5-dimethyl-3-(3-ethoxy-1-methyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline.

<Preparation Example 17>

Production of 5,5-dimethyl-3-(3-ethoxy-1-methyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-2-isoxazoline (present compound No. 3-0012)

0.68 g of m-chloroperbenzoic acid (purity: 70%, 2.76 mmoles) was added, with ice-cooling, to a solution of 0.30 g (0.92 mmoles) of 5,5-dimethyl-3-(3-ethoxy-1-methyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline dissolved in 10 ml of chloroform. The mixture was stirred at room temperature for 5 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water,

followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The resulting crystals were washed with hexane to obtain 0.24 g (yield: 73.0%) of 5,5-dimethyl-3-(3-ethoxy-1-methyl-5-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-2-isoxazoline as white crystals (melting point: 124 to 125°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.50 (2H,s), 4.27 (2H,q), 3.86 (3H,s), 3.04 (2H,s),
1.49 (6H,s), 1.39 (3H,t)

<Preparation Example 18>

Production of 5,5-dimethyl-3-(5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline (present compound No. 3-0027)

19.3 g of sodium hydrosulfide (purity: 70%, 344.6 mmoles) was added to a solution of 21.3 g (120.3 mmoles) of 5,5-dimethyl-3-methylsulfonyl-2-isoxazoline dissolved in 200 ml of N,N-dimethylformamide. The mixture was stirred for 1 hour. Thereto were added 16.7 g (121.0 mmoles) of anhydrous potassium carbonate and 18.6 g (120.7 mmoles) of Rongalit. The resulting mixture was stirred for 2 hours. Thereto was added, with ice-cooling,

31.4 g (120.3 mmols) of 4-bromomethyl-5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazole. The resulting mixture was stirred at room temperature for 2 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 29.0 g (yield: 90.3%) of 5,5-dimethyl-3-(5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline as a yellow oily substance.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.24 (2H,s), 3.90 (3H,s), 2.78 (2H,s), 1.42 (6H,s)

<Preparation Example 19>

Production of 5,5-dimethyl-3-(5-methoxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline (present compound No. 3-0028)

0.77 g (4.0 mmols) of sodium methoxide (a 28% methanol solution) was added to a solution of 0.5 g (1.6 mmols) of 5,5-dimethyl-3-(5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline dissolved in 20 ml of methanol. The mixture was stirred for 4 hours under refluxing, to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by

extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum
5 distillation to remove the solvent contained therein, to obtain 0.5 g (yield: 96.7%) of 5,5-dimethyl-3-(5-methoxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline as a yellow oily substance.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

10 4.26 (2H,s), 4.07 (3H,s), 3.72 (3H,s), 2.80 (2H,s),
1.43 (6H,s)

<Preparation Example 20>

Production of 5,5-dimethyl-3-(5-methoxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-2-
15 isoxazoline (present compound No. 3-0013)

1.3 g of m-chloroperbenzoic acid (purity: 70%, 7.5 mmoles) was added, with ice-cooling, to a solution of 0.5 g (1.5 mmoles) of 5,5-dimethyl-3-(5-methoxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline
20 dissolved in 20 ml of chloroform. The mixture was stirred for 1 hour and then at room temperature for 12 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform.
25 The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution, water and an aqueous sodium

chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The resulting solid was washed with
 5 n-hexane to obtain 0.31 g (yield: 58.2%) of 5,5-dimethyl-3-(5-methoxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-2-isoxazoline as a white powder (melting point: 113.0 to 114.0°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

10 4.60 (2H,s), 4.11 (3H,s), 3.79 (3H,s), 3.10 (2H,s),
 1.51(6H,s)

<Preparation Example 21>

Production of 3-(5-(2-chlorophenoxy)-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-
 15 2-isoxazoline (present compound No. 3-0029)

0.2 g (8.3 mmoles) of sodium hydride (purity: 60%) was added, with ice-cooling, to a solution of 0.44 g (3.4 mmoles) of 2-chlorophenol dissolved in 30 ml of N,N-dimethylformamide. The mixture was stirred for 1 hour.
 20 Thereto was added 0.7 g (2.2 mmoles) of 5,5-dimethyl-3-(5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline. The resulting mixture was stirred at 120 to 130°C for 5 hours to give rise to a reaction. After confirmation of the completion of the
 25 reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium

chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein.

The residue was purified by silica gel column

5 chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 0.63 g (yield: 66.7%) of 3-(5-(2-chlorophenoxy)-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline as a yellow oily substance.

10 <Preparation Example 22>

Production of 3-(5-(2-chlorophenoxy)-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0014)

1.0 g of m-chloroperbenzoic acid (purity: 70%, 5.8
15 mmoles) was added, with ice-cooling, to a solution of 0.63 g (1.5 mmoles) of 3-(5-(2-chlorophenoxy)-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline dissolved in 20 ml of chloroform. The mixture was stirred for 1 hour and then at room
20 temperature for 12 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite
25 solution, an aqueous sodium hydrogencarbonate solution, water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate.

The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The resulting solid was washed with n-hexane to obtain 0.31 g (yield: 45.7%) of 3-(5-(2-chlorophenoxy)-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline as a white powder (melting point: 67.0 to 70.0°C).

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

7.50-6.91 (4H,m), 4.45 (2H,s), 3.71 (3H,s),
3.03 (2H,s), 1.47 (6H,s)

<Preparation Example 23>

Production of 3-(5-cyclopentyloxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0030)

To a solution of 0.43 g (1.6 mmoles) of triphenylphosphine dissolved in 10 ml of benzene were added 0.14 g (1.6 mmoles) of cyclopentanol, 0.5 g (1.6 mmoles) of 5,5-dimethyl-3-(5-hydroxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline and 0.7 g (1.6 mmoles) of diethyl azodicarboxylate (a 40% toluene solution). The mixture was stirred at room temperature for 12 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium

sulfate. The resulting organic layer was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate
5 mixed solvent) to obtain 0.52 g (yield: 85.2%) of 3-(5-cyclopentyloxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline as a colorless transparent oily substance.

<Preparation Example 24>

10 Production of 3-(5-cyclopentyloxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0015)

0.85 g of m-chloroperbenzoic acid (purity: 70%, 4.9 mmoles) was added, with ice-cooling, to a solution of
15 0.52 g (1.4 mmoles) of 3-(5-(cyclopentyloxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline dissolved in 20 ml of chloroform. The mixture was stirred for 1 hour and then at room temperature for 12 hours to give rise to a reaction.
20 After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution,
25 water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum

distillation to remove the solvent contained therein.

The resulting solid was washed with n-hexane to obtain 0.2 g (yield: 35.5%) of 3-(5-cyclopentyloxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-

5 dimethyl-2-isoxazoline as a white powder (melting point: 113.0 to 114.0°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

5.03 (1H,br), 4.60 (2H,s), 3.73 (3H,s), 3.05 (2H,s),
1.88-1.70 (8H,m), 1.50 (6H,s)

10 <Preparation Example 25>

Production of 3-(5-cyano-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline
(present compound No. 3-0031)

0.2 g (4.0 mmoles) of sodium cyanide was added to a
15 solution of 0.5 g (1.6 mmoles) of 5,5-dimethyl-3-(5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-2-isoxazoline dissolved in 30 ml of N,N-dimethylformamide. The mixture was stirred at 40°C for 1
hour to give rise to a reaction. After confirmation of
20 the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting
25 solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 0.9 g of crude 3-(5-cyano-1-methyl-3-trifluoromethyl-1H-pyrazol-4-

ylmethylthio)-5,5-dimethyl-2-isoxazoline as a yellow oily substance.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.30 (2H,s), 4.08 (3H,s), 2.81 (2H,s), 1.43 (6H,s)

5 <Preparation Example 26>

Production of 3-(5-cyano-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0016)

2.1 g of m-chloroperbenzoic acid (purity: 70%, 12.2
10 mmoles) was added, with ice-cooling, to a solution of 0.9 g of 3-(5-cyano-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline (crude compound) dissolved in 50 ml of chloroform. The mixture was stirred for 1 hour and then at room temperature for 12
15 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium
20 hydrogencarbonate solution, water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The resulting solid was washed with
25 n-hexane to obtain 0.43 g (yield: 76.4%) of 3-(5-cyano-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline as a white powder (melting

point: 105.0 to 108.0°C).

^1H -NMR [CDCl_3/TMS , δ (ppm)]:

4.73 (2H,s), 4.16 (3H,s), 3.14 (2H,s), 1.53 (6H,s)

<Preparation Example 27>

- 5 Production of 3-(3,5-dichloro-1-ethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0032)

0.6 g of sodium hydrosulfide (purity: 70%, 10.7 mmoles) was added to a solution of 0.7 g (3.7 mmoles) of
10 5,5-dimethyl-3-ethylsulfonyl-2-isoxazoline dissolved in 30 ml of N,N-dimethylformamide. The mixture was stirred for 1 hour. Thereto were added 0.51 g (3.7 mmoles) of anhydrous potassium carbonate and 0.56 g (3.6 mmoles) of Rongalit. The resulting mixture was stirred for 2 hours.
15 Thereto was added, with ice-cooling, 0.9 g (3.5 mmoles) of 4-bromomethyl-3,5-dichloro-1-ethyl-1H-pyrazole. The resulting mixture was stirred at room temperature for 2 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was
20 poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove
25 the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 0.8 g

(yield: 70.8%) of 3-(3,5-dichloro-1-ethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline as a colorless transparent oily substance.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

5 4.14 (2H,s), 4.14 (2H, q), 2.81 (2H,s), 1.43 (6H,s),
 1.42 (3H,t)

<Preparation Example 28>

Production of 3-(3,5-dichloro-1-ethyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline (present
10 compound No. 3-0017)

2.0 g of m-chloroperbenzoic acid (purity: 70%, 11.6 mmoles) was added, with ice-cooling, to a solution of 0.8 g (2.6 mmoles) of 3-(3,5-dichloro-1-ethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline dissolved in 20
15 ml of chloroform. The mixture was stirred for 1 hour and then at room temperature for 12 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting
20 organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution, water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was
25 subjected to vacuum distillation to remove the solvent contained therein. The resulting solid was washed with n-hexane to obtain 0.41 g (yield: 46.6%) of 3-(3,5-

dichloro-1-ethyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline as a white powder (melting point: 105.0 to 107.0°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

5 4.48 (2H,s), 4.19 (2H,q), 3.05 (2H,s), 1.51 (6H,s),
 1.45 (3H,t)

<Preparation Example 29>

Production of 3-(5-chloro-3-difluoromethyl-1-methyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline
10 (present compound No. 3-0020)

1.2 g of sodium hydrosulfide hydrate (purity: 70%, 15.0 mmoles) was added to a solution of 1.9 g (10.0 mmoles) of 5,5-dimethyl-3-ethylsulfonyl-2-isoxazoline dissolved in 30 ml of N,N-dimethylformamide. The mixture
15 was stirred for 2 hours. Thereto were added 2.1 g (15.0 mmoles) of anhydrous potassium carbonate, 2.3 g (15.0 mmoles) of Rongalit and 2.6 g (10.0 mmoles) of 4-bromomethyl-5-chloro-3-difluoromethyl-1-methyl-1H-pyrazole. The resulting mixture was stirred at room
20 temperature for 15 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then
25 dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified

by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 2.1 g (yield: 68.0%) of 3-(5-chloro-3-difluoromethyl-1-methyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline as
5 a colorless viscous liquid ($n_D^{20} = 1.5183$).

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

6.70 (1H,t, $J=54.2$ Hz), 4.24 (2H,s), 3.86 (3H,s),
2.80 (2H,s), 1.42 (6H,s)

<Preparation Example 30>

10 Production of 3-(5-chloro-3-difluoromethyl-1-methyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0018)

3.6 g of m-chloroperbenzoic acid (purity: 70%, 14.5 mmoles) was added, with ice-cooling, to a solution of 1.8
15 g (5.8 mmoles) of 3-(5-chloro-3-difluoromethyl-1-methyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline dissolved in 15 ml of chloroform. The mixture was stirred at room temperature for 22 hours to give rise to a reaction. After the completion of the reaction, the
20 reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution and an aqueous sodium chloride solution in this order and
25 then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The resulting

crystals were washed with hexane to obtain 1.7 g (yield: 85.9%) of 3-(5-chloro-3-difluoromethyl-1-methyl-1H-pyrazol-4-ylmethylsulfonyl)-5,5-dimethyl-2-isoxazoline as white crystals (melting point: 78 to 79°C).

5 $^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

6.80 (1H, t, $J=54.8$ Hz), 4.60 (2H, s), 3.91 (3H, s),
3.08 (2H, s), 1.51 (6H, s)

<Preparation Example 31>

Production of 5,5-dimethyl-3-(5-methyl-3-trifluoromethylisoxazol-4-ylmethylthio)-2-isoxazoline
10 (present compound No. 4-0003)

0.4 g of sodium hydrosulfide hydrate (purity: 70%, 4.6 mmoles) was added to a solution of 0.4 g (2.3 mmoles) of 5,5-dimethyl-3-methylsulfonyl-2-isoxazoline dissolved
15 in 10 ml of N,N-dimethylformamide. The mixture was stirred for 2 hours. Thereto were added 0.3 g (2.3 mmoles) of potassium carbonate, 0.4 g (2.3 mmoles) of Rongalit and 0.5 g (1.8 mmoles) of 4-bromomethyl-5-methyl-3-trifluoromethylisoxazole. The resulting mixture
20 was stirred at room temperature for 14 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution
25 and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was

purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 0.4 g (yield: 70.0%) of 5,5-dimethyl-3-(5-methyl-3-trifluoromethylisoxazol-4-ylmethylthio)-2-isoxazoline.

5 ^1H -NMR [CDCl_3/TMS , δ (ppm)]:

4.11 (2H,s), 2.77 (2H,s), 2.54 (3H,s), 1.42 (6H,s)

<Preparation Example 32>

Production of 5,5-dimethyl-3-(5-methyl-3-trifluoromethylisoxazol-4-ylmethylsulfonyl)-2-isoxazoline
10 (present compound No. 4-0001)

0.8 g of m-chloroperbenzoic acid (purity: 70%, 3.2 mmoles) was added, with ice-cooling, to a solution of 0.4 g (1.3 mmoles) of 5,5-dimethyl-3-(5-methyl-3-trifluoromethylisoxazol-4-ylmethylthio)-2-isoxazoline
15 dissolved in 10 ml of chloroform. The mixture was stirred at room temperature for 4 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer
20 was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation
25 to remove the solvent contained therein. The resulting crystals were washed with hexane to obtain 0.4 g (yield: 95.0%) of 5,5-dimethyl-3-(5-methyl-3-

trifluoromethylisoxazol-4-ylmethylsulfonyl)-2-isoxazoline
as white crystals (melting point: 135 to 136°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.54 (2H,s), 3.11 (2H,s), 2.61 (3H,s), 1.52 (6H,s)

5 <Preparation Example 33>

Production of [(5-chloro-3-methyl-isothiazol-4-yl)-
methylthio]-5,5-dimethyl-2-isoxazoline (present compound
No. 4-0004)

0.82 g of sodium hydrosulfide (purity: 70%, 10.00
10 mmoles) was added at the room temperature to a solution
of 0.89 g (5.00 mmoles) of 5,5-dimethyl-3-methylsulfonyl-
2-isoxazoline dissolved in 10 ml of N,N-dimethylformamide.
The mixture was stirred for 2 hours. Thereto were added
0.70 g (5.00 mmoles) of anhydrous potassium carbonate,
15 0.78 g (5.00 mmoles) of Rongalit and 0.91 g (5.00 mmoles)
of 5-chloro-4-chloromethyl-3-methylisothiazole. The
resulting mixture was stirred at room temperature
overnight to give rise to a reaction. After confirmation
of the completion of the reaction, the reaction mixture
20 was poured into water, followed by extraction with ethyl
acetate. The resulting organic layer was washed with
water and an aqueous sodium chloride solution and then
dried over anhydrous magnesium sulfate. The resulting
solution was subjected to vacuum distillation to remove
25 the solvent contained therein. The residue was purified
by silica gel column chromatography to obtain 1.38 g
(yield: quantitative) of [(5-chloro-3-methyl-isothiazol-

4-yl)-methylthio]-5,5-dimethyl-2-isoxazoline.

<Preparation Example 34>

Production of [(5-chloro-3-methyl-isothiazol-4-yl)-
methylsulfonyl]-5,5-dimethyl-2-isoxazoline (present
5 compound No. 4-0002)

2.96 g of m-chloroperbenzoic acid (purity: 70%,
12.00 mmoles) was added, with ice-cooling, to a solution
of 1.38 g (5.00 mmoles) of [(5-chloro-3-methyl-
isothiazol-4-yl)-methylthio]-5,5-dimethyl-2-isoxazoline
10 dissolved in 20 ml of chloroform. The mixture was
stirred for 1 hour and then at room temperature for
overnight to give rise to a reaction. After the
completion of the reaction, the reaction mixture was
poured into water, followed by extraction with chloroform.
15 The resulting organic layer was washed with an aqueous
sodium hydrogensulfite solution, an aqueous sodium
hydrogencarbonate solution and an aqueous sodium chloride
solution in this order and then dried over anhydrous
magnesium sulfate. The resulting solution was subjected
20 to vacuum distillation to remove the solvent contained
therein. The residue was purified by silica gel column
chromatography to obtain 0.65 g (yield: 47.0%) of [(5-
chloro-3-methyl-isothiazol-4-yl)-methylsulfonyl]-5,5-
dimethyl-2-isoxazoline as a light yellow powder
25 (melting point: 113 to 114°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

8.89 (1H,s), 4.67 (2H,s), 3.05 (2H,s), 2.59 (3H,s),

1.51 (6H,s)

<Preparation Example 35>

Production of 5,5-dimethyl-3-[2,5-dimethyl-4-(1-methoxyiminoethyl)-thiophen-3-ylmethylthio]-2-isoxazoline
5 (present compound No. 2-0002)

0.57 g (6.8 mmoles) of O-methylhydroxylamine hydrochloride and 0.56 g (6.8 mmoles) of sodium acetate were added to a solution of 1.0 g (3.4 mmoles) of 3-(4-acetyl-2,5-dimethylthiophen-3-ylmethylthio)-5,5-dimethyl-
10 2-isoxazoline dissolved in 50 ml of ethanol. The mixture was stirred for 5 hours under refluxing, to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting
15 organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column
20 chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 0.4 g (36.4%) of 5,5-dimethyl-3-[2,5-dimethyl-4-(1-methoxyiminoethyl)-thiophen-3-ylmethylthio]-2-isoxazoline as a yellow oily substance.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

25 4.21 (2H,s), 3.95 (3H,s), 2.76 (2H,s), 2.38 (3H,s),
2.34 (3H,s), 2.13 (3H,s), 1.42 (6H,s)

<Preparation Example 36>

Production of 5,5-dimethyl-3-[2,5-dimethyl-4-(1-methoxyiminoethyl)-thiophen-3-ylmethylsulfonyl]-2-isoxazoline (present compound No. 2-0001)

0.61 g of m-chloroperbenzoic acid (purity: 70%, 3.5
5 mmoles) was added, with ice-cooling, to a solution of 0.4
g (1.2 mmoles) of 5,5-dimethyl-3-[2,5-dimethyl-4-(1-methoxyiminoethyl)-thiophen-3-ylmethylthio]-2-isoxazoline
dissolved in 30 ml of chloroform. The mixture was
stirred for 1 hour and then at room temperature for 12
10 hours to give rise to a reaction. After confirmation of
the completion of the reaction, the reaction mixture was
poured into water, followed by extraction with chloroform.
The resulting organic layer was washed with an aqueous
sodium hydrogensulfite solution, an aqueous sodium
15 hydrogencarbonate solution, water and an aqueous sodium
chloride solution in this order and then dried over
anhydrous magnesium sulfate. The resulting solution was
subjected to vacuum distillation to remove the solvent
contained therein. The residue was purified by silica
20 gel column chromatography (developing solvent: hexane-
ethyl acetate mixed solvent) to obtain 0.35 g (80%) of
5,5-dimethyl-3-[2,5-dimethyl-4-(1-methoxyiminoethyl)-
thiophen-3-ylmethylsulfonyl]-2-isoxazoline as white
crystals (melting point: 95.0 to 96.0°C).

25 ¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.79 (2H,s), 3.95 (3H,s), 2.93 (2H,s), 2.42 (3H,s),
2.37 (3H,s), 2.17 (3H,s), 1.47 (6H,s)

<Preparation Example 37>

Production of 5,5-dimethyl-3-(4-trifluoromethyl-pyridin-3-ylmethylthio)-2-isoxazoline (present compound No. 7-0003)

5 0.26 g of sodium hydrosulfide (purity: 70%, 4.6 mmoles) was added to a solution of 0.3 g (1.6 mmoles) of 5,5-dimethyl-3-ethylsulfonyl-2-isoxazoline dissolved in 20 ml of N,N-dimethylformamide. The mixture was stirred for 1 hour. Thereto were added 0.22 g (1.6 mmoles) of
10 anhydrous potassium carbonate and 0.25 g (1.6 mmoles) of Rongalit. The resulting mixture was stirred for 2 hours. Thereto was added, with ice-cooling, 0.3 g (1.3 mmoles) of 3-bromomethyl-4-trifluoromethyl-pyridine. The resulting mixture was stirred at room temperature for 2
15 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then
20 dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 0.45 g
25 (yield: 98.9%) of 5,5-dimethyl-3-(4-trifluoromethyl-pyridin-3-ylmethylthio)-2-isoxazoline as a yellow oily substance.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

8.98 (1H,s), 8.70 (1H,d), 7.51 (1H,d), 4.47 (2H,s),
2.79 (2H,s), 1.43 (6H,s)

<Preparation Example 38>

5 Production of 5,5-dimethyl-3-(4-trifluoromethyl-pyridin-3-ylmethylsulfonyl)-2-isoxazoline (present compound No. 7-0001) and 5,5-dimethyl-3-(4-trifluoromethyl-pyridine-N-oxide-3-ylmethylsulfonyl)-2-isoxazoline (present compound No. 7-0002)

10 0.77 g of m-chloroperbenzoic acid (purity: 70%, 4.5 mmoles) was added, with ice-cooling, to a solution of 0.45 g (1.6 mmoles) of 5,5-dimethyl-3-(4-trifluoromethyl-pyridin-3-ylmethylthio)-2-isoxazoline dissolved in 20 ml of chloroform. The mixture was stirred for 1 hour and
15 then at room temperature for 12 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium
20 hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution, water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent
25 contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 0.06 g (yield:

12.0%) of 5,5-dimethyl-3-(4-trifluoromethyl-pyridin-3-ylmethylsulfonyl)-2-isoxazoline as light yellow crystals (melting point: 77.0 to 80.0°C) and 0.12 g (yield: 23.1%) of 5,5-dimethyl-3-(4-trifluoromethyl-pyridin-N-oxide-3-ylmethylsulfonyl)-2-isoxazoline as white crystals (melting point: 114.0 to 116.0°C).

5,5-Dimethyl-3-(4-trifluoromethyl-pyridin-3-ylmethylsulfonyl)-2-isoxazoline

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

10 8.98 (1H,s), 8.84 (1H,d), 7.64 (1H,d), 4.92 (2H,s),
 3.09 (2H,s), 1.52 (6H,s)

5,5-Dimethyl-3-(4-trifluoromethyl-pyridin-N-oxide-3-ylmethylsulfonyl)-2-isoxazoline

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

15 8.50 (1H,s), 8.25 (1H,d), 7.59 (1H,d), 4.81 (2H,s),
 3.12 (2H,s), 1.53 (6H,s)

<Preparation Example 39>

Production of 5,5-dimethyl-[(4-methoxy-6-trifluoromethylpyrimidin-5-yl)-methylthio]-2-isoxazoline
20 (present compound No. 8-0002)

0.32 g of sodium hydrosulfide (purity: 70%, 4.00 mmoles) was added, at room temperature, to a solution of 0.35 g (2.00 mmoles) of 5,5-dimethyl-3-methylsulfonyl-2-isoxazoline dissolved in 10 ml of dimethylformamide. The mixture was stirred for 2 hours. To the reaction mixture were added 0.28 g (2.00 mmoles) of anhydrous potassium carbonate, 0.31 g (2.00 mmoles) of Rongalit and 0.45 g

(2.00 mmoles) of 5-chloromethyl-4-methoxy-6-trifluoromethylpyrimidine. The resulting mixture was stirred at room temperature for 2 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography to obtain 0.55 g (yield: 85.9%) of 5,5-dimethyl-[(4-methoxy-6-trifluoromethylpyrimidin-5-yl)-methylthio]-2-isoxazoline.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

8.81 (1H,s), 4.44 (2H,d), 4.12 (3H,s), 2.81 (2H,s),
1.45 (6H,s)

<Preparation Example 40>

Production of 5,5-dimethyl-[(4-methoxy-6-trifluoromethylpyrimidin-5-yl)-methylsulfonyl]-2-isoxazoline (present compound No. 8-0001)

1.05 g of m-chloroperbenzoic acid (purity: 70%, 4.28 mmoles) was added, with ice-cooling, to a solution of 0.55 g (1.71 mmoles) of 5,5-dimethyl-[(4-methoxy-6-trifluoromethylpyrimidin-5-yl)-methylthio]-2-isoxazoline dissolved in 20 ml of chloroform. The mixture was stirred for 1 hour and then at room temperature for 4

hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium
5 hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained
10 therein. The residue was purified by silica gel column chromatography to obtain 0.45 g (yield: 75.0%) of 5,5-dimethyl-[(4-methoxy-6-trifluoromethylpyrimidin-5-yl)-methylsulfonyl]-2-isoxazoline as white feather-like crystals (melting point: 175 to 176°C).

15 ¹H-NMR [CDCl₃/TMS, δ (ppm)]:

8.89 (1H,s), 5.00 (2H,d), 4.11 (3H,s), 3.11 (2H,s),
1.53 (6H,s)

<Preparation Example 41>

Production of 3-(5,5-dimethyl-2-isoxazolin-3-ylthiomethyl)-2-trifluoromethyl-6,7-dihydro-5H-pyrazolo[5,1-b][1,3]oxazine (present compound No. 3-0033)
20

A solution of 0.82 g (2.3 mmoles) of 3-[5-chloro-1-(3-hydroxypropyl)-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio]-5,5-dimethyl-2-isoxazole dissolved in 5 ml
25 of N,N-dimethylformamide was dropwise added to a suspension of 0.11 g (2.8 mmoles) of sodium hydride in 15 ml of N,N-dimethylformamide. After the completion of the

dropwise addition, the resulting mixture was stirred at room temperature for 30 minutes, then heated to 100°C, and stirred for 1 hour to give rise to a reaction. After confirmation of the completion of the reaction, the
5 reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous citric acid solution and an aqueous sodium chloride solution, and then dried over magnesium sulfate. The resulting solution was subjected
10 to vacuum distillation to obtain 0.77 g (yield: 100%) of 3-(5,5-dimethyl-2-isoxazolin-3-ylthiomethyl)-2-trifluoromethyl-6,7-dihydro-5H-pyrazolo[5,1-b][1,3]oxazine.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

15 4.37 (2H,t), 4.19 (2H,t), 4.15 (2H,s), 2.80 (2H,s),
2.31 (2H,m), 1.42 (6H,s)

<Preparation Example 42>

Production of 3-(5,5-dimethyl-2-isoxazolin-3-ylsulfonylmethyl)-2-trifluoromethyl-6,7-dihydro-5H-
20 pyrazolo[5,1-b][1,3]oxazine (present compound No. 3-0019)

1.25 g of m-chloroperbenzoic acid (purity: 70%, 5.1 mmoles) was added, with ice-cooling, to a solution of 0.77 g (2.3 mmoles) of 3-(6,7-dihydro-3-trifluoromethyl-5H-pyrazolo[5,1-b][1,3]oxazin-4-yl-methylthio)-5,5-
25 dimethyl-2-isoxazoline dissolved in 20 ml of chloroform. The mixture was stirred for 1 hour and then at room temperature for 12 hours to give rise to a reaction.

After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution, water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein.

10 The residue was purified by silica gel column chromatography to obtain 0.36 g (yield: 43%) of 3-(5,5-dimethyl-2-isoxazolin-3-ylsulfonylmethyl)-2-trifluoromethyl-6,7-dihydro-5H-pyrazolo[5,1-b][1,3]oxazine as a white powder (melting point: 151.0 to 152.0°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.47 (2H,s), 4.40 (2H,t), 4.23 (2H,t), 3.09 (2H,s),
2.34 (2H,m), 1.50 (6H,s)

<Preparation Example 43>

20 Production of 3-(5-chloro-1-methyl-3-trifluoromethylpyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0186)

5.59 g of sodium hydrosulfide hydrate (purity: 70%, 69.8 mmoles) was added, at room temperature, to a solution of 6.84 g (35.8 mmoles) of 5,5-dimethyl-3-ethanesulfonyl-2-isoxazoline dissolved in 200 ml of N,N-dimethylformamide. The mixture was stirred for 1 hour.

To the reaction mixture, were added 4.94 g (35.8 mmols) of anhydrous potassium carbonate, 5.51 g (35.8 mmols) of Rongalit and further 9.46 g (34.1 mmols) of 4-bromomethyl-5-chloro-1-methyl-3-trifluoromethyl-1H-pyrazole. The resulting mixture was stirred at room temperature over one night to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography to obtain 8.97 g (yield: 80.3%) of 3-(5-chloro-1-methyl-3-trifluoromethylpyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline.

<Preparation Example 44>

Production of 3-(5-chloro-1-methyl-3-trifluoromethylpyrazol-4-ylmethanesulfonyl)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0039)

16.87 g of m-chloroperbenzoic acid (purity: 70%, 68.4 mmols) was added, with ice-cooling, to a solution of 8.97 g (27.4 mmols) of 3-(5-chloro-1-methyl-3-trifluoromethylpyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline dissolved in 300 ml of chloroform. The mixture was stirred for 1 hour and then at room

temperature over one night to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed
5 with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution, water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove
10 the solvent contained therein. The solid thus obtained was washed with n-hexane to obtain 9.36 g (yield: 95.1%) of 3-(5-chloro-1-methyl-3-trifluoromethylpyrazol-4-ylmethanesulfonyl)-5,5-dimethyl-2-isoxazoline as white powder (melting point: 115.0 to 116.0°C).

15 ¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.63 (2H,s), 3.95 (3H,s), 3.10 (2H,s), 1.52 (6H,s)

<Preparation Example 45>

Production of 3-(5-difluoromethoxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0187)
20

49.4 g (0.88 mol) of powdery potassium hydroxide and 0.94 g (2.9 mmol) of tetra-n-butylammonium bromide were added at room temperature to a solution of 90.3 g (0.29 mole) of 3-(5-hydroxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline
25 dissolved in 1000 ml of tetrahydrofuran. While cooling the system to 20°C or lower, an excess amount of

chlorodifluoromethane was introduced so as to have the system saturated. Thereafter, the reaction mixture was stirred at room temperature for 17 hours. After confirmation of the completion of the reaction, the
5 reaction solution was poured into ice water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and a saturated aqueous sodium chloride solution in this order, and then dried over anhydrous magnesium sulfate. The resulting solution
10 was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel chromatography to obtain 66.6 g (purity: 85%, yield; 54.0%) of 3-(5-difluoromethoxy-1-methyl-3-trifluoro-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-
15 isoxazoline.

<Preparation Example 46>

Production of 3-(5-difluoromethoxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethanesulfonyl)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0188)
20 157.6 g of m-chloroperbenzoic acid (purity: 70%, 0.64 mole) was added, with ice-cooling, to a solution of 56.5 g (0.16 mole) of 3-(5-difluoromethoxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline dissolved in 1000 ml of chloroform. The
25 mixture was stirred at room temperature for 4 hours to give rise to a reaction. After the completion of the reaction, the reaction solution was filtrated to remove

insoluble matters. The filtrate thus obtained was poured into ice water, followed by extraction with chloroform. The resulting organic layer was washed with a 10% aqueous sodium hydroxide solution, water, an aqueous sodium
5 hydrogensulfite solution, and a saturated aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was washed with n-hexane
10 to obtain 52.7 g (yield: 86.0%) of 3-(5-difluoromethoxy-1-methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethanesulfonyl)-5,5-dimethyl-2-isoxazoline as white crystals (melting point: 129.0 to 130.0°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

15 6.83 (1H,t,J=71.9 Hz), 4.60 (2H,s), 3.88 (3H,s),
 3.11 (2H,s), 1.52 (6H,s)

<Preparation Example 47>

Production of 3-(5-difluoromethoxy-1-ethyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio-5,5-dimethyl-2-
20 isoxazoline (present compound No. 3-0189)

30.1 g (536.6 mmol) of powdery potassium hydroxide and 0.5 g of tetra-n-butylammonium bromide were added to a solution of 34.7 g (107.3 mmoles) of 3-(1-ethyl-5-hydroxy-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-
25 dimethyl-2-isoxazoline dissolved in 100 ml of dichloromethane. An excess amount of chlorodifluoromethane was introduced so as to have the

system saturated. Thereafter, the reaction mixture was stirred for 3 hours. After confirmation of the completion of the reaction, the reaction solution was poured into water, followed by extraction with chloroform.

5 The resulting organic layer was washed with water and a saturated aqueous sodium chloride solution in this order, and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was
10 purified by silica gel chromatography to obtain 26.3 g (yield; 65.5%) of 3-(5-difluoromethoxy-1-ethyl-3-trifluoro-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline.

<Preparation Example 48>

15 Production of 3-(5-difluoromethoxy-1-ethyl-3-trifluoromethyl-1H-pyrazol-4-ylmethanesulfonyl)-5,5-dimethyl-2-isoxazoline (present compound No. 3-0190)

30.5 g of m-chloroperbenzoic acid (purity: 70%, 123.9 mmoles) was added, with ice-cooling, to a solution
20 of 18.5 g (49.6 mmoles) of 3-(5-difluoromethoxy-1-ethyl-3-trifluoromethyl-1H-pyrazol-4-ylmethylthio)-5,5-dimethyl-2-isoxazoline dissolved in 200 ml of chloroform. The mixture was stirred at room temperature over one night to give rise to a reaction. After the completion
25 of the reaction, the reaction solution was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium

hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution, water, and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was
5 subjected to vacuum distillation to remove the solvent contained therein. The residue was washed with n-hexane to obtain 19.3 g (yield: 96.0%) of 3-(5-difluoromethoxy-1-ethyl-3-trifluoromethyl-1H-pyrazol-4-ylmethanesulfonyl)-5,5-dimethyl-2-isoxazoline as white
10 powder (melting point: 98 to 100°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

6.83 (1H,t,J=72.0 Hz), 4.60 (2H,s), 4.19 (2H,q),

3.11 (2H,s), 1.52 (6H,s), 1.49 (3H,t)

(Preparation Examples of intermediates)

15 <Reference Example 1>

Production of 3-chloro-5,5-dimethyl-2-isoxazoline

534.0 g (4.0 moles) of N-chlorosuccinimide was gradually added, at 65 to 70°C, to a solution of 182.7 g (2.05 moles) of glyoxylic acid aldoxime dissolved in 2
20 liters of 1,2-dimethoxyethane. The mixture was refluxed for 1 hour with heating. Thereto were added, with ice-cooling, 1,440.0 g (14.4 moles) of potassium hydrogencarbonate and 10 ml of water. Then, 360.0 g (6.4 moles) of 2-methylpropene was added. The resulting
25 mixture was stirred at room temperature for 24 hours to give rise to a reaction. The reaction mixture was poured into water, followed by extraction with diisopropyl ether.

The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 107.7 g (yield: 40.0%) of 3-chloro-5,5-dimethyl-2-isoxazoline as a yellow viscous liquid.

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

2.93 (2H,s), 1.47 (6H,s)

10 <Reference Example 2>

Production of 3-chloro-5-ethyl-5-methyl-2-isoxazoline

61.9 g (463.4 mmol) of N-chlorosuccinimide was gradually added, at 60°C, to a solution of 20.6 g (231.7 mmol) of glyoxylic acid aldoxime dissolved in 500 ml of 1,2-dimethoxyethane. After the addition, the mixture was refluxed for 10 minutes with heating. Thereto were added, with ice-cooling, 50 ml (463.4 mmol) of 2-methyl-1-butene, 98.9 g (1,622 mmol) of potassium hydrogencarbonate and 10 ml of water. The resulting mixture was stirred for 12 hours to give rise to a reaction. The reaction mixture was poured into water, followed by extraction with n-hexane. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 13.9 g (yield: 40.6%) of 3-

chloro-5-ethyl-5-methyl-2-isoxazoline as a yellow viscous liquid.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

2.91 (2H, ABq, J=17.0, Δν=46.1 Hz), 1.73 (2H,q),

5 1.42 (3H,s), 0.96 (3H,t)

<Reference Example 3>

Production of 3-benzylthio-5,5-dimethyl-2-isoxazoline

3.2 g (23.2 mmoles) of anhydrous potassium carbonate and 3.0 g (22.5 mmoles) of 3-chloro-5,5-dimethyl-2-isoxazoline were added, in a nitrogen atmosphere, to a solution of 2.8 g (22.5 mmoles) of benzylmercaptan dissolved in 50 ml of N,N-dimethylformamide. The mixture was stirred at 100°C for 2 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography to obtain 3.1 g (yield: 62.0%) of 3-benzylthio-5,5-dimethyl-2-isoxazoline as a yellow oily substance ($n_D^{20} = 1.5521$).

25 ¹H-NMR [CDCl₃/TMS, δ (ppm)]:

7.24-7.39 (5H,m), 4.26 (2H,s), 2.77 (2H,s),

1.40 (6H,s)

<Reference Example 4>

Production of 3-(2,6-difluorobenzylsulfinyl)-5-ethyl-5-methyl-2-isoxazoline

4.6 g of m-chloroperbenzoic acid (purity: 70%, 18.8
5 mmoles) was added, with ice-cooling, to a solution of 4.1
g (15.0 mmoles) of 3-(2,6-difluorobenzylthio)-5-ethyl-5-
methyl-2-isoxazoline dissolved in 50 ml of chloroform.
The mixture was stirred for 1 hour and then at room
temperature for 12 hours to give rise to a reaction.
10 After the completion of the reaction. the reaction
mixture was poured into water, followed by extraction
with chloroform. The resulting organic layer was washed
with an aqueous sodium hydrogensulfite solution, an
aqueous sodium hydrogencarbonate solution, water and an
15 aqueous sodium chloride solution in this order and then
dried over anhydrous magnesium sulfate. The resulting
solution was subjected to vacuum distillation to remove
the solvent contained therein. The residue was purified
by silica gel column chromatography (developing solvent:
20 hexane-ethyl acetate mixed solvent) to obtain 1.5 g
(yield: 34.8%) of 3-(2,6-difluorobenzylsulfinyl)-5-ethyl-
5-methyl-2-isoxazoline as a white powder (melting point:
30°C or less).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

25 7.39-7.28 (1H,m), 7.03-6.94 (2H,m), 4.38 (2H,s),
3.04 (1H, ABq, J=17.2, Δν=85.7 Hz), 3.12 (1H,s),
1.75 (2H,m), 1.44 (3H,S)+1.41 (3H,s), 0.97 (3H,m)

<Reference Example 5>

Production of 3-(2,6-difluorobenzylsulfonyl)-5-ethyl-5-methyl-2-isoxazoline

1.0 g of m-chloroperbenzoic acid (purity: 70%, 4.1
5 mmoles) was added, with ice-cooling, to a solution of 0.8
g (2.8 mmoles) of 3-(2,6-difluorobenzylsulfinyl)-5-ethyl-
5-methyl-2-isoxazoline dissolved in 50 ml of chloroform.
The mixture was stirred for 1 hour and then at room
temperature for 12 hours to give rise to a reaction.
10 After the completion of the reaction. the reaction
mixture was poured into water, followed by extraction
with chloroform. The resulting organic layer was washed
with an aqueous sodium hydrogensulfite solution, an
aqueous sodium hydrogencarbonate solution, water and an
15 aqueous sodium chloride solution in this order and then
dried over anhydrous magnesium sulfate. The resulting
solution was subjected to vacuum distillation to remove
the solvent contained therein. The residue was purified
by silica gel column chromatography (developing solvent:
20 hexane-ethyl acetate mixed solvent) to obtain 0.6 g
(yield: 75.0%) of 3-(2,6-difluorobenzylsulfonyl)-5-ethyl-
5-methyl-2-isoxazoline as a white powder (melting point:
64 to 65°C).

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

25 7.36-7.46 (1H,m), 6.98-7.04 (2H,m), 4.73 (2H,s),
3.04 (2H, ABq, J=17.2, Δν=51.1 Hz), 1.77 (2H,q),
1.46 (3H,s), 0.97 (3H,t)

<Reference Example 6>

Production of 5,5-dimethyl-3-methylsulfonyl-2-isoxazoline

1.0 kg of an aqueous sodium methanethiolate solution (content: 15%, 2.14 mmoles) was dropwise added, with ice-cooling, to a solution of 143.0 g (1.07 moles) of 3-chloro-5,5-dimethyl-2-isoxazoline dissolved in 500 ml of N,N-dimethylformamide. The mixture was stirred at room temperature for 12 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 115.0 g (yield: 74.1%) of 5,5-dimethyl-3-methylthio-2-isoxazoline. This residue (741.2 mmoles) was dissolved in 1 liter of chloroform. Thereto was added, with ice-cooling, 392.0 g of m-chloroperbenzoic acid (purity: 70%, 1.59 moles). The resulting mixture was stirred for 1 hour and then at room temperature for 12 hours to give rise to a reaction. After the completion of the reaction, the separated m-chloroperbenzoic acid was removed by filtration. The resulting filtrate was washed with an aqueous sodium hydrogensulfite solution, water, an aqueous sodium hydrogencarbonate solution and an aqueous sodium chloride solution in this order and then dried over anhydrous

magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was washed with diisopropyl ether to obtain 77.6 g (yield: 59.1%) of 5,5-dimethyl-3-methylsulfonyl-2-isoxazoline as a white powder (melting point: 82 to 84°C).

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

3.26 (3H,s), 3.12 (2H,s), 1.51 (6H,s)

<Reference Example 7>

10 Production of 5,5-dimethyl-3-ethylthio-2-isoxazoline

1,500 ml of an aqueous solution containing 560.0 g (9.0 moles) of ethyl mercaptan and 360.0 g (9.0 moles) of sodium hydroxide was added to a solution containing 3-chloro-5,5-dimethyl-2-isoxazoline. The mixture was stirred at 60 to 70°C for 16 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 270.0 g of crude 5,5-dimethyl-3-ethylthio-2-isoxazoline as a dark red oily substance.

25 <Reference Example 8>

Production of 5,5-dimethyl-3-ethylsulfonyl-2-isoxazoline

270.0 g (1.7 moles) of crude oily 5,5-dimethyl-3-

ethylthio-2-isoxazoline was dissolved in 1.0 liter of chloroform. Thereto was added, with ice-cooling, 1,050 g of m-chloroperbenzoic acid (purity: 70%, 6.1 moles). The resulting mixture was stirred for 1 hour and then at room temperature for 12 hours to give rise to a reaction. After confirmation of the completion of the reaction, the separated m-chloroperbenzoic acid was removed by filtration. The resulting filtrate was washed with an aqueous sodium hydrogensulfite solution, an aqueous sodium hydrogencarbonate solution, water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was washed with n-hexane to obtain 133.6 g (yield: 65.4%) of 5,5-dimethyl-3-ethylsulfonyl-2-isoxazoline as a white powder.

<Reference Example 9>

Production of 1-phenyl-3-trifluoromethyl-1H-pyrazol-5-ol
20 g (184.9 mmoles) of phenylhydrazine and 4 ml of concentrated hydrochloric acid were added to a solution of 34.1 g (184.9 mmoles) of ethyl trifluoroacetoacetate dissolved in 500 ml of ethanol. The mixture was refluxed for 1 hour with heating, to give rise to a reaction. After the completion of the reaction, the reaction mixture was subjected to vacuum distillation to remove the most part of the solvent contained therein. The residue was mixed with water to precipitate crystals.

The crystals were collected by filtration, washed with water until the filtrate became neutral, and dried to obtain 37.1 g (yield: 87.9%) of 1-phenyl-3-trifluoromethyl-1H-pyrazol-5-ol as ocherous crystals.

5 ^1H -NMR [CDCl_3/TMS , δ (ppm)]:

7.68-7.41 (5H,m), 5.86 (1H,s), 3.71 (1H,s)

<Reference Example 10>

Production of 5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazole-4-carboaldehyde

10 33.6 g (219.1 mmol) of phosphorus oxychloride was added to 7.7 g (105.2 mmol) of N,N-dimethylformamide with ice-cooling. Thereto was added, at room temperature, 20 g (87.7 mmol) of 1-phenyl-3-trifluoromethyl-1H-pyrazol-5-ol. The resulting mixture was refluxed for 1
15 hour with heating, to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water with ice-cooling, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium hydrogencarbonate
20 solution and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column
25 chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 19.1 g (yield: 79.1%) of 5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazole-4-

carboaldehyde as white crystals.

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

10.06 (1H,s), 7.57 (5H,s)

<Reference Example 11>

5 Production of (5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-yl)-methanol

A solution of 0.21 g (5.5 mmoles) of lithium aluminum hydride dissolved in 70 ml of THF was cooled to -30°C. Thereto was gradually added a solution of 3 g (10.9 mmoles) of 5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazole-4-carboaldehyde dissolved in 30 ml of tetrahydrofuran. The resulting mixture was stirred at -30°C for 30 minutes to give rise to a reaction. After the completion of the reaction, ethyl acetate was added, followed by stirring. Then, water was added, followed by stirring for a while. The reaction mixture was filtered under vacuum. The filtrate was extracted with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 3.0 g (yield: 99.9%) of (5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-yl)-methanol as white crystals.

25 $^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

7.54-7.51 (5H,m), 4.71 (2H,d) 1.79 (1H,b)

<Reference Example 12>

Production of 4-bromomethyl-5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazole

A solution of 3.0 g (10.9 mmol) of (5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-yl)-methanol dissolved in 60 ml of diethyl ether was cooled to -10°C. Thereto was added 1.0 g (3.8 mmol) of phosphorus tribromide. The mixture was stirred at room temperature for 1 hour to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 3.6 g (yield: 95.8%) of 4-bromomethyl-5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazole as white crystals.

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

7.58-7.48 (5H, m), 4.48 (2H, s)

<Reference Example 13>

Production of 5-fluoro-1-phenyl-3-trifluoromethyl-1H-pyrazole-4-carbaldehyde

10.5 g (180.2 mmol) of potassium fluoride was added to a solution of 33.0 g (120.1 mmol) of 5-chloro-1-phenyl-3-trifluoromethyl-1H-pyrazole-4-carbaldehyde dissolved in 500 ml of dimethyl sulfoxide. The mixture was stirred at 100°C for 2 hours to give rise to a

reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 26.5 g (yield: 85.0%) of 5-fluoro-1-phenyl-3-trifluoromethyl-1H-pyrazole-4-carboaldehyde.

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

9.96 (1H,s), 7.68-7.51 (5H,m)

<Reference Example 14>

15 Production of (5-fluoro-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-yl)-methanol

To a solution of 1.6 g (41.0 mmoles) of sodium borohydride dissolved in 300 ml of methanol was added, with ice-cooling, a solution of 26.5 g (102.5 mmoles) of 5-fluoro-1-phenyl-3-trifluoromethyl-1H-pyrazole-4-carboaldehyde dissolved in 200 ml of methanol. The resulting mixture was stirred at 0°C for 30 minutes to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate.

The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 28.5 g (yield: 100%) of (5-fluoro-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-yl)-methanol.

5 ^1H -NMR [CDCl_3/TMS , δ (ppm)]:

7.65-7.41 (5H,m), 4.68 (2H,d), 1.73 (1H,t)

<Reference Example 15>

Production of 4-bromomethyl-5-fluoro-1-phenyl-3-trifluoromethyl-1H-pyrazole

10 A solution of 27.5 g (105.7 mmoles) of (5-fluoro-1-phenyl-3-trifluoromethyl-1H-pyrazol-4-yl)-methanol dissolved in 300 ml of diethyl ether was cooled to 0°C. Thereto was added 10.0 g (37.0 mmoles) of phosphorus tribromide. The mixture was stirred at room temperature
15 for 2 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with diethyl ether. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over
20 anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 30.3 g (yield: 88.8%) of 4-bromomethyl-5-fluoro-1-phenyl-3-trifluoromethyl-1H-pyrazole.

25 ^1H -NMR [CDCl_3/TMS , δ (ppm)]:

7.66-7.42 (5H,m), 4.44 (2H,s)

<Reference Example 16>

Production of 1-tert-butyl-3-trifluoromethyl-1H-pyrazol-5-ol

373.8 g (3.0 moles) of tert-butylhydrazine hydrochloride and 50 ml of concentrated hydrochloric acid were added to a solution of 552.3 g (3.0 moles) of ethyl trifluoroacetoacetate dissolved in 1,500 ml of ethanol. The mixture was refluxed for 2 days with heating, to give rise to a reaction. After the completion of the reaction, the reaction mixture was subjected to vacuum distillation to remove the most part of the solvent contained therein. The residue was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was washed with n-hexane to obtain 369.0 g (yield: 59.1%) of 1-tert-butyl-3-trifluoromethyl-1H-pyrazol-5-ol as a white powder.

<Reference Example 17>

Production of 1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazole-4-carboaldehyde

462.0 g (3.0 moles) of phosphorus oxychloride was added to 87.7 g (1.2 moles) of N,N-dimethylformamide with ice-cooling. Thereto was added, at room temperature, 208.2 g (1.0 moles) of 1-tert-butyl-3-trifluoromethyl-1H-pyrazol-5-ol. The resulting mixture was refluxed for 10

hours with heating, to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with water, a 5% aqueous sodium hydroxide solution and water in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 131.5 g (yield: 21.7%) of 1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazole-4-carboaldehyde as white crystals.

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

9.97 (1H, d), 1.76 (9H, s)

<Reference Example 18>

Production of (1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazol-4-yl)-methanol

A solution of 39.9 g (156.9 mmol) of (1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazole-4-carboaldehyde dissolved in 300 ml of methanol was cooled to 0°C. Thereto was gradually added 6.5 g (172.6 mmol) of sodium borohydride. The mixture was stirred at room temperature for 3 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was

washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 37.7 g (yield: 93.6%) of (1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazol-4-yl)-methanol.

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

4.60 (2H,d), 1.72 (9H,s), 1.58 (1H,t)

<Reference Example 19>

10 Production of 4-bromomethyl-1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazole

A solution of 9.2 g (35.7 mmol) of (1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazol-4-yl)-methanol dissolved in 100 ml of diethyl ether was cooled to -10°C . Thereto was added 11.6 g (42.9 mmol) of phosphorus tribromide. The mixture was stirred at room temperature overnight to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with diethyl ether. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 10.0 g (yield: 87.3%) of 4-bromomethyl-1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazole.

<Reference Example 20>

Production of (1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazol-4-yl)-methanethiol

43.5 g (136.1 mmoles) of 4-bromomethyl-1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazole was added to a solution of 21.8 g of sodium hydrosulfide hydrate (purity: 70%, 272.2 mmoles) dissolved in 300 ml of N,N-dimethylformamide. The mixture was stirred at room temperature overnight to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with diethyl ether. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 32.3 g (yield: 87.0%) of (1-tert-butyl-5-chloro-3-trifluoromethyl-1H-pyrazol-4-yl)-methanethiol.

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

3.65 (2H,d), 1.90 (1H,t), 1.70 (9H,s)

<Reference Example 21>

Production of 1-tert-butyl-5-methoxy-3-trifluoromethyl-1H-pyrazole

15.0 g (108.4 mmoles) of anhydrous potassium carbonate and 19.3 g (135.5 mmoles) of methyl iodide were added, at room temperature, to a solution of 18.8 g (90.3 mmoles) of 1-tert-butyl-3-trifluoromethyl-1H-pyrazol-5-ol dissolved in 100 ml of N,N-dimethylformamide. The

mixture was stirred for 15 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with diethyl ether. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 20.0 g (yield: 99.8%) of 1-tert-butyl-5-methoxy-3-trifluoromethyl-1H-pyrazole.

<Reference Example 22>

Production of 1-tert-butyl-4-chloromethyl-5-methoxy-3-trifluoromethyl-1H-pyrazole

5.4 g of paraformaldehyde (180.2 mmols in terms of formaldehyde) and 20 ml of concentrated hydrochloric acid were added to a solution of 20.0 g (90.1 mmols) of 1-tert-butyl-5-methoxy-3-trifluoromethyl-1H-pyrazole dissolved in 90 ml of acetic acid. The mixture was stirred at 60°C for 30 minutes to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with diisopropyl ether. The resulting organic layer was washed with water and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 21.7 g (yield: 89.0%) of 1-tert-butyl-4-chloromethyl-5-methoxy-3-trifluoromethyl-1H-pyrazole.

<Reference Example 23>

Production of 3-methoxy-1-methyl-5-trifluoromethyl-1H-pyrazole

10.0 g (72.3 mmoles) of anhydrous potassium
5 carbonate and 12.8 g (90.3 mmoles) of methyl iodide were added, at room temperature, to a solution of 10.0 g (60.2 mmoles) of 3-hydroxy-1-methyl-5-trifluoromethyl-1H-pyrazole dissolved in 50 ml of N,N-dimethylformamide. The mixture was stirred for 15 hours to give rise to a
10 reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with diethyl ether. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over
15 anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 9.8 g (yield: 90.7%) of 3-methoxy-1-methyl-5-trifluoromethyl-1H-pyrazole.

<Reference Example 24>

20 Production of 4-chloromethyl-3-methoxy-1-methyl-5-trifluoromethyl-1H-pyrazole

0.45 g of paraformaldehyde (15.0 mmoles in terms of formaldehyde) and 5 ml of concentrated hydrochloric acid were added to a solution of 1.00 g (5.6 mmoles) of 3-
25 methoxy-1-methyl-5-trifluoromethyl-1H-pyrazole dissolved in 25 ml of acetic acid. The mixture was stirred at 80°C for 2 hours to give rise to a reaction. After the

completion of the reaction, the reaction mixture was poured into water and neutralized with potassium carbonate, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 0.83 g (yield: 65.0%) of 4-chloromethyl-3-methoxy-1-methyl-5-trifluoromethyl-1H-pyrazole.

<Reference Example 25>

Production of 5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazole-4-carboaldehyde

42.0 g (711.9 mmols) of potassium fluoride was added to a solution of 60.4 g (282.7 mmols) of 5-chloro-1-methyl-3-trifluoromethyl-1H-pyrazole-4-carboaldehyde dissolved in 700 ml of dimethyl sulfoxide. The mixture was stirred at 120 to 140°C for 5 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column

chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 36.8 g (yield: 66.0%) of 5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazole-4-carboaldehyde.

5 <Reference Example 26>

Production of (5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazol-4-yl)-methanol

To a solution of 3.9 g (102.6 mmoles) of sodium borohydride dissolved in 500 ml of methanol was added,
10 with ice-cooling, a solution of 36.8 g (187.6 mmoles) of 5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazole-4-carboaldehyde dissolved in 200 ml of methanol. The resulting mixture was stirred at 0°C for 30 minutes to give rise to a reaction. After confirmation of the
15 completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting
20 solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 35.4 g (yield: 95.4%) of (5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazol-4-yl)-methanol.

<Reference Example 27>

25 Production of 4-bromomethyl-5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazole

A solution of 35.4 g (178.7 mmoles) of 5-fluoro-1-

methyl-3-trifluoromethyl-1H-pyrazole-4-methanol dissolved in 500 ml of diethyl ether was cooled to -30°C . Thereto was added 54.0 g (199.5 mmoles) of phosphorus tribromide. The mixture was stirred at room temperature for 12 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with diethyl ether. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 31.4 g (yield: 80.8%) of 4-bromomethyl-5-fluoro-1-methyl-3-trifluoromethyl-1H-pyrazole.

<Reference Example 28>

Production of (ethoxycarbonyl)malondialdehyde

12.6 g of sodium hydride (purity: 60%, 525.0 mmoles) was washed with diethyl ether by decantation several times and then made into a solution in 500 ml of diethyl ether. Thereto were added, in a nitrogen current at 0 to 10°C , 194 g (2.6 moles) of ethyl formate and 50 g (262.0 mmoles) of ethyl 3,3-diethoxy-propionate. The resulting mixture was stirred at room temperature for 15 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by washing with diethyl ether. The resulting aqueous layer was allowed to have a pH of 1

with hydrochloric acid, followed by extraction with dichloromethane. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution
5 was subjected to vacuum distillation to remove the solvent contained therein, to obtain 37.6 g (yield: 100%) of crude (ethoxycarbonyl)malondialdehyde as a dark red oily substance.

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

10 9.09 (2H,s), 5.26 (1H,s), 4.27 (2H,q), 1.28 (3H,t)

<Reference Example 29>

Production of ethyl 1H-pyrazole-4-carboxylate

6.2 g (193 mmoles) of hydrazine was added, with ice-cooling, to a solution of 27.6 g (192 mmoles) of
15 (ethoxycarbonyl)malondialdehyde dissolved in 150 ml of ethanol. The mixture was stirred at room temperature for 17 hours to give rise to a reaction. The reaction mixture was subjected to vacuum distillation to remove the ethanol contained therein. The residue was purified
20 by silica gel column chromatography (developing solvent: dichloromethane-ethyl acetate mixed solvent) to obtain 19.4 g (72.4%) of ethyl 1H-pyrazole-4-carboxylate as yellow crystals.

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

25 8.08 (2H,s), 5.30 (1H,s), 4.31 (2H,q), 1.36 (3H,t)

<Reference Example 30>

Production of ethyl 1-ethyl-1H-pyrazole-4-carboxylate

3.7 g (26.8 mmol) of anhydrous potassium carbonate and 4.2 g (26.6 mmol) of ethyl iodide were added to a solution of 1.5 g (10.7 mmol) of ethyl 1H-pyrazole-4-carboxylate dissolved in 50 ml of N,N-dimethylformamide. The mixture was stirred at room temperature for 20 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 1.6 g (yield: 88.9%) of ethyl 1-ethyl-1H-pyrazole-4-carboxylate as a yellow oily substance.

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

7.90 (2H,s), 4.28 (2H,q), 4.18 (2H,q), 1.51 (3H,t),
1.35 (3H,t)

<Reference Example 31>

Production of ethyl 3,5-dichloro-1-ethyl-1H-pyrazole-4-carboxylate

In a glass sealed tube were placed 1.6 g (9.5 mmol) of ethyl 1-ethyl-1H-pyrazole-4-carboxylate and 5.1 g (38.3 mmol) of N-chlorosuccinimide. There were allowed to react at 160°C for 6 hours. After the

completion of the reaction, the reaction mixture was cooled to room temperature, washed with carbon tetrachloride and chloroform, and filtered under vacuum. The resulting filtrate (organic layer) was washed with
5 water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent:
10 hexane-ethyl acetate mixed solvent) to obtain 1.0 g (yield: 44.2%) of ethyl 3,5-dichloro-1-ethyl-1H-pyrazole-4-carboxylate as a yellow oily substance.

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

4.36 (2H,q), 4.21 (2H,q), 1.44 (3H,t), 1.38 (3H,t)

15 <Reference Example 32>

Production of (3,5-dichloro-1-ethyl-1H-pyrazol-4-yl)methanol

A solution of 0.16 g (4.2 mmoles) of lithium aluminum hydride dissolved in 70 ml of tetrahydrofuran
20 was cooled to -50°C . Thereto was gradually added dropwise a solution of 1.0 g (4.2 mmoles) of ethyl 3,5-dichloro-1-ethyl-1H-pyrazole-4-carboxylate dissolved in 30 ml of tetrahydrofuran. The mixture was stirred at -50°C for 3 hours to give rise to a reaction. After
25 confirmation of the completion of the reaction, ethyl acetate was added, followed by stirring for a while. Water was added, followed by stirring for a while. The

resulting mixture was filtered under vacuum. The filtrate was extracted with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 0.82 g (yield: 100%) of (3,5-dichloro-1-ethyl-1H-pyrazol-4-yl)methanol as a brown oily substance.

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

4.52 (2H,s), 4.16 (2H,q), 1.43 (3H,t)

<Reference Example 33>

Production of 4-bromomethyl-3,5-dichloro-1-ethyl-1H-pyrazole

A solution of 0.82 g (4.2 mmol) of (3,5-dichloro-1-ethyl-1H-pyrazol-4-yl)methanol dissolved in 50 ml of diethyl ether was cooled to -30°C . Thereto was added 1.3 g (4.8 mmol) of phosphorus tribromide. The mixture was stirred at room temperature for 12 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 0.9 g (yield: 81.8%) of 4-bromomethyl-3,5-dichloro-1-ethyl-1H-pyrazole as a yellow oily substance.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.33 (2H,s), 4.13 (2H,q), 1.43 (3H,t)

<Reference Example 34>

Production of 3-difluoromethyl-1-methyl-1H-pyrazol-5-ol

- 5 8.3 g (180.6 mmoles) of methylhydrazine and 5 ml of concentrated hydrochloric acid were added to a solution of 30.0 g (180.6 mmoles) of ethyl difluoroacetoacetate dissolved in 200 ml of ethanol. The mixture was refluxed for 2 days with heating, to give rise to a reaction.
- 10 After the completion of the reaction, the reaction mixture was subjected to vacuum distillation to remove the most part of the solvent contained therein. The residue was poured into water. The mixture was allowed to have a pH of 4 using citric acid and extracted with
- 15 ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein.
- 20 The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 8.9 g (yield: 33.3%) of 3-difluoromethyl-1-methyl-1H-pyrazol-5-ol.

<Reference Example 35>

- 25 Production of 5-chloro-3-difluoromethyl-1-methyl-1H-pyrazole-4-carboaldehyde

41.6 g (270.1 mmoles) of phosphorus oxychloride was

added, with ice-cooling, to 7.9 g (108.0 mmol) of N,N-dimethylformamide. Thereto was added, at room temperature, 8.0 g (54.0 mmol) of 3-difluoromethyl-1-methyl-1H-pyrazol-5-ol. The mixture was refluxed for 4 hours with heating, to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with water, a 5% aqueous sodium hydroxide solution and water in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 7.7 g (yield: 73.3%) of 5-chloro-3-difluoromethyl-1-methyl-1H-pyrazole-4-carbaldehyde as white crystals.

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

9.96 (1H,s), 6.90 (1H,t, $J=53.6$ Hz), 3.93 (3H,s)

<Reference Example 36>

Production of (5-chloro-3-difluoromethyl-1-methyl-1H-pyrazol-4-yl)-methanol

A solution of 7.2 g (37.0 mmol) of 5-chloro-3-difluoromethyl-1-methyl-1H-pyrazole-4-carbaldehyde dissolved in 100 ml of methanol was cooled to 0°C . Thereto was gradually added 2.1 g (55.5 mmol) of sodium borohydride. The mixture was stirred at room temperature for 3 hours to give rise to a reaction. After the

completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 3.8 g (yield: 52.1%) of (5-chloro-3-difluoromethyl-1-methyl-1H-pyrazol-4-yl)-methanol.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

6.70 (1H, t, J=40.8 Hz), 4.63 (2H, s), 3.86 (3H, s),
1.79 (1H, br)

<Reference Example 37>

Production of 4-bromomethyl-5-chloro-3-difluoromethyl-1-methyl-1H-pyrazole

A solution of 2.0 g (10.0 mmol) of (5-chloro-3-difluoromethyl-1-methyl-1H-pyrazol-4-yl)-methanol dissolved in 50 ml of diethyl ether was cooled to -10°C. Thereto was added 1.0 g (3.5 mmol) of phosphorus tribromide. The mixture was stirred at room temperature overnight to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into ice water, followed by extraction with diethyl ether. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the

solvent contained therein, to obtain 2.6 g (yield: 100.0%) of 4-bromomethyl-5-chloro-3-difluoromethyl-1-methyl-1H-pyrazole.

<Reference Example 38>

5 Production of trifluoroacetaldehyde oxime etherate

24.1 g (347.0 mmoles) of hydroxylamine hydrochloride and 160 ml of water were added to a solution of 50.0 g (347.0 mmoles) of trifluoroacetaldehyde hemiethyl acetal dissolved in 80 ml of methanol. Thereto was dropwise
10 added, with ice-cooling, 80.0 g of a 50% aqueous sodium hydroxide solution (1.7 moles). After the completion of the dropwise addition, the resulting mixture was stirred at room temperature for 6 hours to give rise to a reaction. After the completion of the reaction, 10%
15 hydrochloric acid was added for pH adjustment to 6. The resulting mixture was extracted with diethyl ether. The extract was subjected to vacuum distillation to remove the solvent contained therein. The residue was subjected to distillation to obtain 24.7 g (yield: 38.0%) of
20 trifluoroacetaldehyde oxime etherate.

<Reference Example 39>

Production of trifluoroacetohydroximoyl bromide etherate

A solution of 38.8 g (218.0 mmoles) of N-bromosuccinimide dissolved in 125 ml of N,N-
25 dimethylformamide was added, with ice-cooling, to a solution of 24.7 g (131.7 mmoles) of trifluoroacetaldehyde oxime etherate dissolved in 50 ml

of N,N-dimethylformamide. The mixture was stirred at room temperature for 3 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with diethyl ether. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was subjected to distillation to obtain 33.3 g (yield: 95.0%) of trifluoroacetoxyhydroxymoyl bromide etherate as a brown oily substance.

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

9.30 (1H, s)

<Reference Example 40>

Production of 4-ethoxycarbonyl-5-methyl-3-trifluoromethylisoxazole

2.8 g (51.3 mmol) of sodium methoxide was added to a solution of 6.7 g (51.3 mmol) of ethyl acetoacetate dissolved in 80 ml of methanol. Thereto was added, with ice-cooling, a solution of 5.0 g (18.8 mmol) of trifluorohydroxymoyl bromide etherate dissolved in 20 ml of methanol. The resulting mixture was stirred at room temperature for 3 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was subjected to vacuum distillation to remove the solvent contained therein. Water was added to the

residue, followed by extraction with chloroform. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum
5 distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 2.9 g (yield: 69.0%) of 4-ethoxycarbonyl-5-methyl-3-trifluoromethylisoxazole as a
10 colorless oily substance.

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

4.36 (2H,q), 2.77 (3H,s), 1.37 (3H,t)

<Reference Example 41>

Production of (5-methyl-3-trifluoromethylisoxazol-4-yl)-
15 methanol

A solution of 0.16 g (4.2 mmol) of lithium aluminum hydride dissolved in 15 ml of THF was cooled to 0°C. Thereto was gradually added a solution of 0.93 g (4.2 mmol) of 4-ethoxycarbonyl-5-methyl-3-
20 trifluoromethylisoxazole dissolved in 15 ml of THF. The mixture was stirred at 0°C for 1 hour to give rise to a reaction. After the completion of the reaction, ethyl acetate was added, followed by stirring for a while. Water was added, followed by stirring for a while. The
25 reaction mixture was filtered under vacuum. The filtrate was extracted with diethyl ether. The resulting organic layer was washed with an aqueous sodium chloride solution

and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 0.5 g (yield: 60.0%) of (5-methyl-3-trifluoromethylisoxazol-4-yl)-methanol.

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

4.60 (2H,d), 2.54 (3H,s), 1.66 (1H, br)

<Reference Example 42>

Production of 4-bromomethyl-5-methyl-3-

trifluoromethylisoxazole

A solution of 0.45 g (2.5 mmoles) of (5-methyl-3-trifluoromethylisoxazol-4-yl)-methanol dissolved in 10 ml of diethyl ether was cooled to 0°C. Thereto was added 0.2 g (8.9 mmoles) of phosphorus tribromide. The mixture was stirred at room temperature for 1 hour to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with diethyl ether. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 0.5 g (yield: 74.0%) of 4-bromomethyl-5-methyl-3-trifluoromethylisoxazole.

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

4.31 (2H,d), 2.51 (3H,s)

<Reference Example 43>

Production of (5-chloro-3-methyl-isothiazol-4-yl)-methanol

A solution of 2.06 g (10.0 mmoles) of ethyl 5-chloro-3-methyl-isothiazole-4-carboxylate dissolved in 10 ml of THF was dropwise added, at -30°C, to a solution of 0.42 g (11.0 mmoles) of lithium aluminum hydride dissolved in 10 ml of THF. The mixture was stirred at the same temperature for 1 hour to give rise to a reaction. After confirmation of the completion of the reaction, ethyl acetate was added to the reaction mixture. The resulting mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography to obtain 1.50 g (yield: 91.5%) of (5-chloro-3-methyl-isothiazol-4-yl)-methanol.

<Reference Example 44>

Production of 4-chloromethyl-5-chloro-3-methylisothiazole

3.26 g (27.44 mmoles) of thionyl chloride was added, at room temperature, to a solution of 1.50 g (9.15 mmoles) of (5-chloro-3-methyl-isothiazol-4-yl)-methanol dissolved in 10 ml of chloroform. The mixture was stirred for 3 hours to give rise to a reaction. After confirmation of the completion of the reaction, the

reaction mixture was subjected to vacuum distillation to remove the solvent contained therein, to obtain 1.67 g (yield: quantitative) of 4-chloromethyl-5-chloro-3-methylisothiazole.

5 <Reference Example 45>

Production of methyl 4-trifluoromethylnicotinate

6.7 g (48.6 mmoles) of anhydrous potassium carbonate and 6.9 g (48.6 mmoles) of methyl iodide were added to a solution of 4.6 g (24.1 mmoles) of 4-
10 trifluoromethylnicotinic acid dissolved in 70 ml of N,N-dimethylformamide. The mixture was stirred at room temperature for 12 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by
15 extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein.
20 The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 2.77 g (yield: 56.1%) of methyl 4-trifluoromethylnicotinate as a yellow oily substance.

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

25 9.11 (1H,s), 8.92 (1H,d), 7.64 (1H,d), 3.99 (3H,s)

<Reference Example 46>

Production of (4-trifluoromethylpyridin-3-yl)-methanol

A solution of 0.37 g (9.7 mmol) of lithium aluminum hydride dissolved in 100 ml of THF was cooled to -50°C. Thereto was gradually added dropwise a solution of 2.0 g (9.8 mmol) of methyl 4-trifluoromethylnicotinate dissolved in 30 ml of THF. The mixture was stirred at -50°C for 3 hours to give rise to a reaction. After confirmation of the completion of the reaction, ethyl acetate was added, followed by stirring for a while. Water was added, followed by stirring for a while. The reaction mixture was filtered under vacuum. The filtrate was extracted with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 0.6 g (yield: 35.3%) of (4-trifluoromethylpyridin-3-yl)-methanol as a yellow oily substance.

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

9.00 (1H,s), 8.73 (1H,d), 7.51 (1H,d), 4.95 (2H,s)

<Reference Example 47>

Production of 3-bromomethyl-4-trifluoromethylpyridine

A solution of 0.6 g (3.4 mmol) of (4-trifluoromethylpyridin-3-yl)-methanol dissolved in 50 ml of diethyl ether was cooled to -30°C. Thereto was added

1.4 g (5.2 mmoles) of phosphorus tribromide. The mixture was stirred at room temperature for 12 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 0.61 g (yield: 75.3%) of 3-bromomethyl-4-trifluoromethylpyridine as a yellow oily substance.

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

8.88 (1H,s), 8.73 (1H,d), 7.54 (1H,d), 4.63 (2H,s)

<Reference Example 48>

15 Production of 5-bromo-4-hydroxy-6-trifluoromethylpyrimidine

77.5 g (945.0 mmoles) of anhydrous sodium acetate was added, at room temperature, to a solution of 49.2 g (300.0 mmoles) of 4-hydroxy-6-trifluoromethylpyrimidine dissolved in 600 ml of acetic acid. Thereto was gradually added 50.3 g (315 mmoles) of bromine at 45°C. The resulting mixture was stirred at the same temperature for 3 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was subjected to vacuum distillation to remove the solvent contained therein. The residue was poured into water, followed by extraction with ethyl

acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum
5 distillation to remove the solvent contained therein. The residue was washed with n-hexane to obtain 38.9 g (yield: 53.4%) of 5-bromo-4-hydroxy-6-trifluoromethylpyrimidine.

<Reference Example 49>

10 Production of 5-bromo-4-chloro-6-trifluoromethylpyrimidine

24.3 g (100.0 mmoles) of 5-bromo-4-hydroxy-6-trifluoromethylpyrimidine was suspended in 18.5 g (120.0 mmoles) of phosphorus oxychloride. The mixture was
15 stirred at 100°C for 2 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water gradually, followed by extraction with chloroform. The resulting organic layer was washed with water and an aqueous sodium
20 chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography to obtain 21.5 g (yield: 82.4%)
25 of 5-bromo-4-chloro-6-trifluoromethylpyrimidine.

<Reference Example 50>

Production of 5-bromo-4-methoxy-6-

trifluoromethylpyrimidine

16.7 ml of sodium methoxide (a 28% methanol solution, 86.4 mmoles) was added, at room temperature, to a solution of 21.5 g (82.2 mmoles) of 5-bromo-4-chloro-6-trifluoromethylpyrimidine dissolved in 100 ml of methanol. The mixture was stirred to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was subjected to vacuum distillation to remove the solvent contained therein. The residue was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was washed with n-hexane to obtain 19.2 g (yield: 91.0%) of 5-bromo-4-methoxy-6-trifluoromethylpyrimidine.

<Reference Example 51)

Production of 5-bromo-4-ethoxy-6-trifluoromethylpyrimidine

0.94 g (13.77 mmoles) of sodium ethoxide was added, at room temperature, to a solution of 3.00 g (11.48 mmoles) of 5-bromo-4-chloro-6-trifluoromethylpyrimidine dissolved in 50 ml of ethanol. The mixture was stirred to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was subjected to vacuum distillation to remove the solvent

contained therein. The residue was poured into water, followed by extraction with chloroform. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over
5 anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography to obtain 2.44 g (yield: 82.9%) of 5-bromo-4-ethoxy-6-trifluoromethylpyrimidine.

10 <Reference Example 52>

Production of 4-methoxy-6-trifluoromethylpyrimidine-5-carboaldehyde

30.0 ml of n-butyllithium (a 1.6 moles/liter n-hexane solution, 48.0 mmoles) was gradually added, at -65
15 to -60°C, to a solution of 10.3 g (40.0 mmoles) of 5-bromo-4-methoxy-6-trifluoromethylpyrimidine dissolved in 100 ml of tetrahydrofuran. The mixture was stirred for 30 minutes. Thereto was added 3.6 g (48.0 mmoles) of ethyl formate at the same temperature. The resulting
20 mixture was stirred at the same temperature for 3 hours to give rise to a reaction. The reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this
25 order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein.

The residue was purified by silica gel column chromatography to obtain 1.3 g (yield: 15.8%) of 4-methoxy-6-trifluoromethylpyrimidine-5-carboaldehyde.

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

5 10.41 (1H,q), 8.91 (1H,s), 4.18 (3H,s)

<Reference Example 53>

Production of 4-ethoxy-6-trifluoromethylpyrimidine-5-carboaldehyde

A solution of 5.76 g (21.3 mmoles) of 5-bromo-4-ethoxy-6-trifluoromethylpyrimidine dissolved in 250 ml of THF was cooled to -78°C . Thereto was dropwise added 22.6 ml of n-butyllithium (a 1.6 moles/liter n-hexane solution, 36.1 mmoles) . The mixture was stirred for 40 minutes. Thereto was added 2.7 g (45.1 mmoles) of methyl formate.

15 The resulting mixture was stirred for 1.5 hours to give rise to a reaction. After the completion of the reaction, an aqueous ammonium chloride solution was added. The mixture was extracted with diethyl ether. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate.

20 The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate mixed solvent) to obtain 3.82 g (yield: 81.6%) of 4-ethoxy-6-trifluoromethylpyrimidine-5-carboaldehyde.

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

10.41 (1H,s), 8.95 (1H,s), 4.63 (2H,q), 1.48 (3H,t)

<Reference Example 54>

Production of (4-methoxy-6-trifluoromethylpyrimidin-5-yl)-methanol

5 0.24 g (6.3 mmoles) of sodium borohydride was gradually added, at room temperature, to a solution of 1.3 g (6.3 mmoles) of 4-methoxy-6-trifluoromethylpyrimidine-5-carboaldehyde dissolved in 30 ml of methanol. The mixture was stirred for 3 hours to
10 give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this
15 order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography to obtain 0.42 g (yield: 32.1%) of (4-methoxy-6-trifluoromethylpyrimidin-5-yl)-methanol
20

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

8.93 (1H,s), 4.81 (2H,s), 4.13 (3H,s), 2.26 (1H,br)

<Reference Example 55>

Production of (4-ethoxy-6-trifluoromethylpyrimidin-5-yl)-
25 methanol

A solution of 3.82 g (17.2 mmoles) of 4-ethoxy-6-trifluoromethylpyrimidine-5-carboaldehyde dissolved in 50

ml of methanol was added, with ice-cooling, to a solution of 1.7 g (45.7 mmoles) of sodium borohydride dissolved in 50 ml of methanol. The mixture was stirred at 0°C for 1 hour to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 3.77 g (yield: 97.8%) of (4-ethoxy-6-trifluoromethylpyrimidin-5-yl)-methanol

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

8.80 (1H,s), 4.81 (2H,s), 4.59 (2H,q), 2.28 (1H,b),
1.48 (3H,t)

<Reference Example 56>

Production of 5-chloromethyl-4-methoxy-6-trifluoromethylpyrimidine

1.19 g (10.1 mmoles) of thionyl chloride was added, at room temperature, to a solution of 0.42 g (2.02 mmoles) of (4-methoxy-6-trifluoromethylpyrimidin-5-yl)-methanol. The mixture was stirred for 3 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was subjected to vacuum distillation to remove the solvent contained therein, to obtain 0.45 g (yield: quantitative) of 5-chloromethyl-4-methoxy-6-trifluoromethylpyrimidine.

<Reference Example 57>

Production of 5-bromomethyl-4-ethoxy-6-trifluoromethylpyrimidine

A solution of 3.77 g (17.0 mmoles) of (4-ethoxy-6-trifluoromethylpyrimidin-5-yl)-methanol dissolved in 50 ml of diethyl ether was cooled to 0°C. Thereto was added 2.0 g (7.2 mmoles) of phosphorus tribromide. The mixture was stirred at room temperature for 1 hour. The resulting salt was dissolved using methanol. The resulting mixture was stirred for 1 hour to give rise to a reaction. The reaction mixture was poured into water, followed by extraction with diethyl ether. The resulting organic layer was washed with an aqueous sodium chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain crude 5-bromomethyl-4-ethoxy-6-trifluoromethylpyrimidine.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

8.79 (1H,s), 4.61 (2H,q), 4.55 (2H,s), 1.49 (3H,t)

<Reference Example 58>

Production of 4-methoxy-6-trifluoromethylpyrimidine-5-carboaldehyde

30.0 ml (48.0 mmol) of n-butyllithium (1.6 mol/l n-hexane solution) was slowly added to 100 ml of tetrahydrofuran solution containing 10.3 g (40.0 mmol) of 5-bromo-4-methoxy-6-trifluoromethylpyrimidine at -65 to -

60°C, and the resultant mixture was stirred at 30 minutes. Further, 3.6 g (48.0 mmol) of ethyl formate was added thereto at the same temperature, and the mixture was stirred for 3 hours at the same temperature. The resultant reaction solution was poured into water and was extracted with ethyl acetate. The organic phase thus obtained was washed with water and an aqueous sodium chloride solution in this order, and then dried over anhydrous magnesium sulfate. The resultant solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography to obtain 1.3 g (yield: 15.8%) of 4-methoxy-6-trifluoromethylpyrimidine-5-carboaldehyde.

¹H-NMR [CDCl₃/TMS δ (ppm)] 10.41(1H,q) 8.98(1H,s),
4.18(3H,s)

<Reference Example 59>

Production of (2-chloro-4-methylpyridin-3-yl)methanol

A solution of 1.9 g (10.0 mmoles) of methyl 2-chloro-4-methylnicotinate dissolved in 5.0 ml of THF was gradually added, at -65 to -60°C, to a suspension of 0.4 g (10.0 mmoles) of lithium aluminum hydride in 30 ml of tetrahydrofuran. The mixture was stirred for 30 minutes and at -20°C for 1 hour to give rise to a reaction. The reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over

anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography to obtain 0.6 g (yield: 38.2%) of (2-chloro-4-methylpyridin-3-yl)methanol.

$^1\text{H-NMR}$ [CDCl_3/TMS , δ (ppm)]:

8.19 (1H,d), 7.08 (1H,d), 4.85 (2H,s), 2.49 (3H,s)

<Reference Example 60>

Production of 3-acetyl-4-chloromethyl-2,5-dichlorothiophene

33 ml of titanium tetrachloride (a 2 moles/liter dichloromethane solution, 66.0 mmoles) was dropwise added, at 10°C with ice-cooling, to a solution of 5.0 g (32.4 mmoles) of 3-acetyl-2,5-dichlorothiophene dissolved in 26 ml (323.0 mmoles) of chloromethyl methyl ether. The mixture was stirred at room temperature for 2 hours to give rise to a reaction. After the completion of the reaction, the reaction mixture was poured into ice water, followed by extraction with chloroform. The resulting organic layer was washed with sodium bicarbonate, water and an aqueous sodium chloride solution in this order and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane/ethyl acetate = 9/1) to obtain 2.6 g (yield: 39.7%) of 3-acetyl-4-chloromethyl-2,5-

dichlorothiophene as yellow crystals.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

4.70 (2H,s), 2.56 (3H,s), 2.54 (3H,s), 2.39 (3H,s)

<Reference Example 61>

5 Production of 3-bromo-2-bromomethylbenzofuran

2.7 g (15.3 mmoles) of N-bromosuccinimide and 0.4 g (2.7 mmoles) of azobisisobutyronitrile were added to a solution of 2.8 g (13.3 mmoles) of 3-bromo-2-methylbenzofuran dissolved in 30 ml of monochlorobenzene.

10 The mixture was stirred at 80°C for 30 minutes to give rise to a reaction. After confirmation of the disappearance of the raw materials, the reaction mixture was cooled to room temperature. The insolubles were removed by filtration. The filtrate was subjected to
15 vacuum distillation to remove the solvent contained therein. The residue was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium chloride solution in this order and then dried over
20 anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein, to obtain 3.0 g (yield: 79.0%) of 3-bromo-2-bromomethylbenzofuran.

<Reference Example 62>

25 Production of ethyl 1-difluoromethyl-1H-pyrazole-4-carboxylate

6.0 g (43.5 mmoles) of anhydrous potassium carbonate

was added to a solution of 3.0 g (21.4 mmoles) of ethyl 1H-pyrazole-4-carboxylate dissolved in 100 ml of N,N-dimethylformamide. Thereinto was blown chlorodifluoromethane. The resulting mixture was stirred
5 at 130 to 140°C for 3 hours to give rise to a reaction. After confirmation of the completion of the reaction, the reaction mixture was poured into water, followed by extraction with ethyl acetate. The resulting organic layer was washed with water and an aqueous sodium
10 chloride solution and then dried over anhydrous magnesium sulfate. The resulting solution was subjected to vacuum distillation to remove the solvent contained therein. The residue was purified by silica gel column chromatography (developing solvent: hexane-ethyl acetate
15 mixed solvent) to obtain 1.67 g (yield: 41.0%) of ethyl 1-difluoromethyl-1H-pyrazole-4-carboxylate as a colorless transparent oily substance.

¹H-NMR [CDCl₃/TMS, δ (ppm)]:

8.32 (1H,s), 8.04 (1H,s), 7.20 (1H,t), 4.32 (2H,q),
20 1.37 (3H,t)

Now, Working Examples are illustrated below. In the Examples, "part" means "part by weight".

<Formulation 1> Wettable powder

5 Parts of compound No. 3-0002 and 40 parts of
25 cyanazine were mixed with 0.5 part of polyoxyethylene octylphenyl ether, 0.5 part of a sodium salt of an alkaylnaphthalenesulfonic acid-formalin condensate, 12

parts of diatomaceous earth and 42 parts of clay. The mixture was mixed and pulverized to obtain a wettable powder.

The amount of the herbicidal composition of the present invention varies depending on a mixing ratio, weather conditions, a form of formulation, an application time, an application method, an application place, a target weed, a target crop or the like, but is usually from 50 to 1500 g per hectare as a total amount of active ingredients. In a case of an emulsion, a wettable powder, a suspension or the like, a predetermined amount is diluted with water in an amount of from 100 to 1000 liters per hectare to be applied.

The effect of the herbicidal composition of the present invention is illustrated by the following Application Examples.

<Application Example 1> Test for herbicidal effect by upland field soil treatment

An upland field soil was filled in a plastic pot of 11 cm each of length, width and depth. Seeds of corn, green foxtail (*Setaria viridis*) and common lambsquarters (*Chenopodium album* L.) were sowed, followed by covering with the same soil. The wettable powder produced in accordance with Formulation 1 was weighed so as to provide active ingredients in a predetermined amount, and was diluted with water and sprayed uniformly on the soil surface using a small sprayer, in an amount of 100 liters

per 10 ares. Then, growing was made in a greenhouse, and the herbicidal effect of each wettable powder was examined at the 30th day from the treatment in accordance with the standard shown in Table 15. The results are shown in Tables 16 and 17.

Table 15

Index	Herbicidal effect (extent of growth inhibition) or phytotoxicity
10	Herbicidal effect of growth inhibition or phytotoxicity of 100%
9	Herbicidal effect or phytotoxicity of from 90 to 99%
8	Herbicidal effect or phytotoxicity of from 80 to 89%
7	Herbicidal effect or phytotoxicity of from 70 to 79%
6	Herbicidal effect or phytotoxicity of from 60 to 69%
5	Herbicidal effect or phytotoxicity of from 50 to 59%
4	Herbicidal effect or phytotoxicity of from 40 to 49%
3	Herbicidal effect or phytotoxicity of from 30 to 39%
2	Herbicidal effect or phytotoxicity of from 20 to 29%
1	Herbicidal effect or phytotoxicity of from 10 to 19%
0	Herbicidal effect or phytotoxicity of from 0 to 9%

Table 16

Compound	Amounts of active ingredient (g a.i./ha)	Herbicidal effect		
		Corn	Green foxtail (Sataria viridis)	Common lambsquarter (Chenopodium album L.)
3-0002	32	0	5	2
Cyanazine	500	0	1	2
3-0002 + Cyanazine	32+500	0	10	7
3-0004	32	0	3	2
Cyanazine	500	0	1	2
3-0004 + Cyanazine	32+500	0	9	7
3-0005	32	0	2	2
Cyanazine	500	0	1	2
3-0005 + Cyanazine	32+500	0	9	6
3-0010	16	0	2	2
Cyanazine	500	0	1	2
3-0010 + Cyanazine	16+500	0	10	8
3-0011	16	0	3	2
Cyanazine	500	0	1	2
3-0011 + Cyanazine	16+500	0	10	8
3-0012	16	0	2	1
Cyanazine	500	0	1	2
3-0012 + Cyanazine	16+500	0	9	7
3-0013	16	0	2	2
Cyanazine	500	0	1	2
3-0013 + Cyanazine	16+500	0	9	7
3-0014	32	0	2	2
Cyanazine	500	0	1	2
3-0014 + Cyanazine	32+500	0	9	8
3-0015	32	0	3	3
Cyanazine	500	0	1	2
3-0015 + Cyanazine	32+500	0	9	8
3-0016	16	0	3	3
Cyanazine	500	0	1	2
3-0016 + Cyanazine	16+500	0	10	9
3-0017	16	0	3	3
Cyanazine	500	0	1	2
3-0017 + Cyanazine	16+500	0	9	9
3-0018	16	0	3	3
Cyanazine	500	0	1	2
3-0018 + Cyanazine	16+500	0	10	9
4-0001	32	0	2	1
Cyanazine	500	0	1	2
4-0001 + Cyanazine	32+500	0	8	7

Table 17

Compound	Amounts of active ingredient (g a.i./ha)	Herbicidal effect		
		Corn	Green foxtail (Sataria viridis)	Common lambsquarter (Chenopodium album L.)
4-0002	32	0	3	2
Cyanazine	500	0	1	2
4-0002 + Cyanazine	32+500	0	9	7
2-0001	32	0	2	1
Cyanazine	500	0	1	2
2-0001 + Cyanazine	32+500	0	8	7
8-0001	16	0	2	1
Cyanazine	500	0	1	2
8-0001 + Cyanazine	16+500	0	9	7
3-0039	16	0	3	3
Cyanazine	500	0	1	2
3-0039 + Cyanazine	16+500	0	10	9
3-0188	16	0	3	2
Cyanazine	500	0	1	2
3-0188 + Cyanazine	16+500	0	10	9
3-0190	16	0	3	1
Cyanazine	500	0	1	2
3-0190 + Cyanazine	16+500	0	10	9

<Application Example 2> Test for herbicidal effect

by upland field soil treatment

- 5 An upland field soil was filled in a plastic pot of 11 cm each of length, width and depth. Seeds of corn and velvetleaf (*Abutilon theophrasti* Medic) were sowed, followed by covering with the same soil. The wettable powder produced in accordance with Formulation 1 was
- 10 weighed so as to provide active ingredients in a predetermined amount, and was diluted with water and sprayed uniformly on the soil surface using a small

sprayer, in an amount of 100 liters per 10 ares. Then, growing was made in a greenhouse, and the herbicidal effect of each wettable powder was examined at the 30th day from the treatment in accordance with the standard shown in Table 15. The results are shown in Table 18.

Table 18

Compound	Amounts of active ingredient (g a.i./ha)	Corn	Velvetleaf (Abutilon theophrasti Medic)
Compound 3-188	16	0	2
Compound 3-188	32	0	2
Atrazine	125	0	1
Compound-3188 Atrazine	16-125	0	7
Compound 3-188 Atrazine	32-125	0	10

INDUSTRIAL APPLICABILITY

The herbicidal composition of the present invention, which comprises a compound of the formula (I) and at least one compound selected from Group A, achieves not a simple total herbicidal activity but a synergistic herbicidal effect. Thus, the composition of the present invention achieves an excellent herbicidal effect at a small dose to various weeds growing on an upland field in a wide term range of from before germination to growing season, such as gramineous weeds including barnyardgrass (*Echinochloa crusgalli* var. *crus-galli*), crabgrass (*Digitaria ciliaris*), green foxtail (*Setaria viridis*),

annual bluegrass (*Poa annua*), johnsongrass (*Sorghum halepense* Pers.), blackgrass (*Alopecurus myosuroides*), wild oats (*Avena fatua*), and the like, broad leaf weeds including pale persicaria (*Polygonum lapathifolia*),

5 slender amaranth (*Amaranthus viridis*), common lambsquarters (*Chenopodium album* L.), common chickweed (*Stellaria media* Villars), velvetleaf (*Abutilon theophrasti* Medic), prickly sida (*Sida spinosa* L.), Hemp sesbania (*Sesbania exaltata* Cory), common ragweed

10 (*Ambrosia artemisiifolia*), morningglory, and the like, and annual and perennial sedge weeds including purple nutsedge (*Cyperus rotundus* L.), yellow nutsedge (*Cyperus esculentus* L.), hime-kugu (*Cyperus brevifolius* H.), sedge weed (*Cyperus microiria* Steud), rice flatsedge (*Cyperus*

15 *iria* L.), and the like.

Further, the composition of the present invention achieves an excellent herbicidal effect at a small dose to various weeds growing on a paddy field in a wide term range of from before germination to growing season, such

20 as annual weeds including watergrass (*Echinochloa oryzicola*), smallflower umbrella plant (*Cyperus difformis*), konagi (*Monochoria vaginalis*), aze-na (*Lindernia procumbens*), and the like and perennial weeds including mizu-gayatsuri (*Cyperus serotinus*), kuroguwai

25 (*Eleocharis kuroguwai*), inu-hotaru-i (*Scirpus juncooides*), and the like.

On the other hand, the herbicidal composition of the

present invention is safe to aimed crop, particularly
safe to rice, wheat, barley, corn, grain sorghum,
soybeans, cotton, sugar beet, turf, fruit trees, and the
like.